



UNIVERSITY  
OF TASMANIA

Evaluating the contribution of tonoplast and  
plasma membrane transporters in salinity  
tissue tolerance in barley

Getnet Dino Adem, B.Sc. (Crop Sciences);

M.Sc. (Crop Production); M.Sc. (Plant Molecular Biology)

This thesis is presented for the degree of

Doctor of Philosophy

at the University of Tasmania

Faculty of Science, Engineering and Technology

School of Land and Food

August 2015

# Declaration

This thesis contains no material which has been accepted for the award of any other degree or diploma in any tertiary institution and to the best of my knowledge contains no material previously published or written by any other person, except where due reference is made in the text of this thesis.

Getnet Dino Adem

This Thesis may be made available for loan and limited copying in accordance with the Copyright Act 1968.

Getnet Dino Adem

University of Tasmania

School of Land and Food

Hobart

August 2015

Dedicated to beloved families and friends

---

## Acknowledgment

First and foremost, I would like to thank Prof. Sergey Shabala for understanding my capability to undertake research at PhD level. I am also grateful to him for materializing my longtime dream. I also thank him for leading this PhD project with full deliberation, readiness to be contacted and have discussion at any time even if when having lunch. Secondly, I would like to pass my profound gratitude collectively for all my supervisors, Prof. Sergey Shabala, Dr. Stuart Roy, A/Prof Meixue Zhou and A/Prof John Bowman for their knowledgeable, skillful and studious assistance throughout all my PhD study. I am also indebted to Dr. Lana Shabala for patiently assisting in MIFE experiments and also in every other issue in the lab. I am also grateful for all the staff members and students of Australian Centre for Plant Functional Genomics (ACPFG), in particular for salt research group.

I also extend thanks for the help rendered at the CSL Molecular Genetics lab at the University of Tasmania, particularly Mr. Adam Smolenski and Ms. Sharee McCammon. I would like to thank Dr. Jim Weller and Mrs. Michelle Lang for providing their quarantine approved premise as well. I am also grateful for Mr. Phil Andrew's help with greenhouse facility and growth condition issues. I also thank all the lab friends at the University of Tasmania, SLAF, plant physiology group under Prof. Sergey Shabala and other fellow school friends. I also would like to thank the University of Tasmania, Graduate Research Office for awarding me scholarship to undertake my PhD study. Last but not least, I thank all my family, Dino, Herya, Kamil, Sanya, Kelifa (Late brother praying from above), Zuriash, Sofia, Razika, my wife Menderin, Eyobed and Fikir for their moral, psychological and spiritual support in times of success and hardship in life.

---

## Abstract

Salinity is one of the major abiotic stresses affecting the world food supply. Salt affected soil has area coverage of 950 million ha which accounts for 10% of the land surface of the globe. Fifty percent of the irrigated land (230 million ha) is affected by salinity costing the world food production \$US27 billion per annum. In Australia alone, 67% of the agriculture land is affected by transient salinity costing the Australian economy \$AUD 1330 million per annum. In comparison to other strategies, the use of salt tolerant plants is cost effective and sustainable way of controlling salinity.

While numerous attempts have been made to develop salt tolerant varieties over the last few decades, most of the efforts were focused on tackling individual components or genes contributing to overall salt tolerance. As a result, the progress in the field was much slower than expected and we still lack truly salt tolerant varieties in the farmers' field. Salinity tolerance is a multi-faceted physiological trait, and all the beneficial effects of improving a function of one gene/mechanism may be lost or overturned by plethora of other contributing factors. This calls for a need of "pyramiding" several key traits in one ideotype. Before this can be implemented into the practice, the essentiality and a relative contribution of various traits should be quantified, for each particular species. The aim of this PhD study was to identifying the major contributing mechanisms to salt tolerance in barley. The major focus was on the following aspects: (1) essentiality of transcriptional vs post translational factors in mediating plant adaptive responses to salinity; (2) quantifying the relative contribution of osmo- and tissue-tolerance mechanisms in barley; (3) identifying components involved in  $\text{Na}^+$  sequestration in vacuole; and (4) revealing the role of  $\text{H}^+$ -ATPase in vacuolar ion sequestration and overall salinity stress tolerance.

In the first part of this PhD study, the relative contribution of ionic, osmotic and oxidative stresses to the overall salinity tolerance in barley was studied, both at the whole

plant and cellular level. In addition, the gene expression profile of key genes in ionic and oxidative homeostasis (*NHX*, *RBOH*, *SOD*, *AHA* and *GORK*) as a way of comparing the contribution of transcriptional and post translational factors for salinity tolerance was also investigated. The major findings can be summarized to two main points. These are: (i) tissue tolerance is the dominant component in which root  $K^+$  retention and lower sensitivity to stress induced hydroxyl radical production are the main ones; (ii) responses at the post-translational level is more important than at the transcriptional level for understanding the mechanisms for salt tolerance. Overall, for better tissue tolerance, sodium sequestration,  $K^+$  retention and resistance to oxidative stress are important salt tolerance mechanisms. It is concluded that every crop improvement programs for salinity stress tolerance should take into consideration all this components.

In the second part of this PhD study, we showed that unlike a number of reports for salt-sensitive “salt excluder” species (such as *Arabidopsis* or rice), expressing *AtNHX1* in barley had no beneficial effect on plant performance under saline conditions. *AtNHX1* *Arabidopsis* tonoplast  $Na^+/H^+$  exchanger was expressed in barley (*Hordeum vulgare* L., cv. Golden Promise) and the plants grown under saline growth condition. The transgenic plants were compared with null segregant for biomass, water content, gas exchange, and  $Na^+$  and  $K^+$  content of the leaf. The transgenic barley plants expressing *AtNHX1* has not shown significant difference from the null segregant for any of the trait at least under our experimental conditions. The lack of phenotype in barley which adapts “salt including” strategy was explained by one or more of the following: (i) low level of activity of vacuolar  $H^+$ -PPiase and vacuolar  $H^+$ -ATPase causing poor proton gradient; (ii) the lack of controlling passive leak of sodium via  $Na^+$  permeable slow activating and fast activating channels in the vacuole; (iii) insufficient ATP pool to assist the  $H^+$  pumping activity; (iv) the *AtNHX1* protein may not be folded properly, inactive or mis-targeted.

In the last part of this PhD study, improvement in salinity stress tolerance was obtained in barley crops by expressing V-ATPase subunit C gene. Most of the reports until now have not shown improvement at the grain yield level and also have not explained the physiological mechanisms behind. Also, no previous attempt to express the V-ATPase subunit C was done in barley. Accordingly, we expressed *AtVHA-C* gene in barley and grown under 300mM NaCl. The barley plants expressing the gene were compared with wild type plants for dry biomass, leaf pigment content, stomatal conductance, grain yield, and leaf Na<sup>+</sup> and K<sup>+</sup> content. The transgenic barley plant expressing *AtVHA-C* have shown smaller reduction in biomass and grain yield compared to wild type plants. The beneficial effect in the *AtVHA-C* expressing plant was due to better maintenance of stomatal conductance resulted from the accumulation of Na<sup>+</sup> and K<sup>+</sup> in the leaf that lead to osmotic adjustment and less reliance on the *de novo* synthesis of organic osmolytes.

To recap, salinity tolerance is a multigenic physiological trait involving various mechanisms that may differ between species. For barley, the dominant mechanism appears to be the tissue tolerance. This includes better K<sup>+</sup> retention in the root; reduced tissue sensitivity to oxidative stress; and more efficient sodium sequestration in the leaf. We have shown that for a better sodium sequestration all the components such as proton pumping for generating proton gradient in the tonoplast membrane, control of the back leak of sodium through Na<sup>+</sup> permeable SV and FV channels, and ensuring properly folded, active and correctly targeted NHX protein to the tonoplast all should be considered and modified as one set, to achieve a positive outcome and improve crop salinity tolerance. Finally, we have also shown the importance of vacuolar H<sup>+</sup>-ATPase towards salinity tolerance by the contribution of vacuolar H<sup>+</sup>-ATPase for the accumulation of Na<sup>+</sup> and K<sup>+</sup> in the leaf where they contribute towards osmotic adjustment, and hence, conservation of energy otherwise spent for organic osmolyte production.

---

## **Publications arising from this thesis**

- (1) Adem GD., Roy S., Zhou M., Bowman J. and Shabala S. (2014) Evaluating contribution of ionic, osmotic and oxidative stress components towards salinity tolerance in barley. *BMC Plant Biology* 14 (1), 113. (Chapter 3)
- (2) Adem GD., Roy S., Plett D., Zhou M., Bowman J. and Shabala S. (2015) Expressing *AtNHX1* in barley (*Hordeum vulgare* L.) does not improve plant performance under saline conditions. *Plant Growth Regul*, 1-9. DOI 10.1007/s10725-015-0063-9 (Chapter 4)
- (3) Adem GD., Roy S. Zhou M., Bowman J. and Shabala S. (2015) Expressing Arabidopsis thaliana V-ATPase Subunit C in barley (*Hordeum vulgare* L.) improves plant performance under saline condition by enabling better osmotic adjustment (submitted to *Plant Biotechnology Journal*). (Chapter 5)
- (4) Adem GD., Bose J., Zhou M., and Shabla S. (2015) Targeting vacuolar sodium sequestration in plant breeding for salinity tolerance. In H. Wani and A. Hossain (eds.) Managing salinity tolerance in plants: Molecular and Genomic perspectives (pp. 35-50), Taylor & Francis Catalogue # K23522 ISBN: 978-1-4822-4513-4. (Chapter 2)



---

## List of Abbreviations

AHA- *Arabidopsis* plasma membrane proton ATPase

AtHKT 1;1- *Arabidopsis thaliana* high affinity K<sup>+</sup> transporter 1;1

AtNHX1- *Arabidopsis thaliana* tonoplast sodium/proton exchanger 1

AtNHX2- *Arabidopsis thaliana* tonoplast sodium/proton exchanger 2

AtNHX3- *Arabidopsis thaliana* tonoplast sodium/proton exchanger 3

AtNHX4- *Arabidopsis thaliana* tonoplast sodium/proton exchanger 4

AtVHA-C- *Arabidopsis thaliana* vacuolar proton ATPase subunit C

AUD- Australian dollar

AVP- *Arabidopsis* vacuolar pyrophosphatase

betB- betaine aldehyde dehydrogenase

BSM- Basic salt medium

BYDV- Barley yellow dwarf virus

CaCl<sub>2</sub>- Calcium chloride

CBL10- Calcineurin binding like 10

CDH/betA- Choline dehydrogenase

CIPK 24- CBL interacting protein kinase 24

CodA- Choline oxidase

DW- Dry weight

FV- Tonoplast fast activating Na<sup>+</sup> permeable K<sup>+</sup> channel

FW-fresh weight

GAPDH2- Gyceraldehyde-3-phosphate dehydrogenase 2

GhNHX1- *Gossypium hirsutum* tonoplast sodium/proton antiporter1

GORK- Gated outward rectified K<sup>+</sup> channel

Gs- Gas exchange/ stomatal conductance

- HAK5- Potassium high affinity uptake transporter 5
- HKT- High affinity  $K^+$  transporter
- HvHKT 2;1- *Hordeum vulgare* high affinity  $K^+$  transporter 2;1
- HvNHX1- *Hordeum vulgare* tonoplast sodium/proton exchanger 1
- HvNHX2- *Hordeum vulgare* tonoplast sodium/proton exchanger 2
- HVP10- *Hordeum vulgare* Pyrophosphatase 10
- Hv-PMHATPase- *Hordeum vulgare* plasma membrane  $H^+$  ATPase
- Hv-VRT2- *Hordeum vulgare* vernalization gene 2
- KCl- Potassium chloride
- KOR- Potassium outward rectified channel
- LeHAK 5- *Lycopersicon esculentum* high affinity  $K^+$  uptake transporter 5
- LeNHX2- *Lycopersicon esculentum* tonoplast sodium/proton exchanger 2
- LIX- Liquid Ion Exchanger
- MATE- Multi drug and toxic extrusion gene
- NaCl- Sodium chloride
- NaClO- Sodium hypochlorite
- NADPH oxidase- Nicotine adenine dinucleotide phosphate oxidase
- Nax2- Sodium exclusion locus
- NHX- Tonoplast sodium/proton exchanger/antiporter
- NHX5- tonoplast sodium/proton exchanger 5
- NHX6- tonoplast sodium/proton exchanger 6
- NSCC- Non selective cation channel
- $O_2^-$ - Superoxide
- PCD- Programmed cell death
- RBoHF- Respiratory burst oxidase subunit F

- ROS- Reactive oxygen species
- RT-PCR- Reverse transcriptase polymerase chain reaction
- RWC- Relative water content
- SbMATE- Sorghum bicolor multi drug and toxic extrusion gene
- SOD- Superoxide dismutase
- SOS1- Salt overly sensitive 1
- SOS2- Salt overly sensitive 2
- SOS3- salt overly sensitive 3
- SsNHX1- *Suaeda salsa* tonoplast sodium/proton exchanger
- SV- Tonoplast slow activating Na<sup>+</sup> permeable K<sup>+</sup> channel
- TGN-Trans-golgi network
- ThNHX1- *Thellinguella halophila* tonoplast sodium/proton antiporter1
- TmHKT1;5-A- *Triticum monococcum* high affinity K<sup>+</sup> transporter 1;5-A
- TPC-Two pore potassium channel
- USDA- United states department of Agriculture
- USD- United States dollar
- V-ATPase- Vacuolar proton ATPase
- VHA- Vacuolar proton ATPase
- VMA- Yeast vacuolar proton ATPase
- V-PPiase- Vacuolar proton pyrophosphatase

## Table of Contents

Acknowledgment.....	iv
Abstract.....	v
Publications arising from this thesis.....	viii
List of Abbreviations.....	ix
Table of Contents .....	xii
List of Figures.....	xvi
List of Tables.....	xix
Chapter 1    General Introduction.....	1
1.1 Salinity's Menace to Agricultural Productivity.....	1
1.2 Mechanisms of plant salt tolerance .....	3
1.2.1 Osmotic adjustment .....	3
1.2.2 Sodium exclusion.....	4
1.2.3 Sodium sequestration .....	6
1.2.4 Potassium retention.....	7
1.2.5 Reactive oxygen species detoxification .....	8
1.3 Research aims.....	8
Chapter 2    Targeting vacuolar sodium sequestration in plant breeding for salinity tolerance (literature review) .....	10
Abstract.....	10
Introduction .....	10
2.1 Essentiality of vacuolar Na <sup>+</sup> sequestration for plant salinity tolerance .....	11
2.2 Molecular and physiological mechanisms for Na <sup>+</sup> loading into vacuole .....	13
2.2.1 Molecular identity of NHX proteins .....	13
2.2.2 NHX proteins and their role in salt tolerance .....	14
2.3 Regulation of NHX proteins.....	15
2.4 Selectivity of NHX proteins and its impact on the activity of K <sup>+</sup> homeostasis.....	17
2.5 Fuelling NHX activity .....	19
2.5.1 Essentiality of H <sup>+</sup> -pumping for the sequestration process .....	19
2.5.2 Regulation of vacuolar H <sup>+</sup> -ATPases under saline conditions.....	20
2.5.3 Regulation of vacuolar H <sup>+</sup> -PPases under saline conditions .....	21
2.6 Retaining Na <sup>+</sup> in the vacuole .....	23
2.6.1 Pathways of Na <sup>+</sup> back-leak into the cytosol .....	23
2.6.2 The molecular identity and regulation of SV and FV vacuolar channels .....	24
2.7 The role of SV and FV channels in plant salinity tolerance.....	25
2.8 Targeting vacuolar sequestration in breeding programs .....	27

2.9 Conclusions .....	30
Chapter 3    Evaluating contribution of ionic, osmotic and oxidative stress components towards salinity tolerance in barley .....	31
Abstract.....	31
3.1 Background.....	32
3.2 Methods .....	36
3.2.1 Plant materials and growth conditions .....	36
3.2.2 Whole-plant agronomical and physiological characteristics .....	38
3.2.3 Tissue sap ion content .....	38
3.2.4 Non-invasive ion flux measurements.....	38
3.2.5 Gene Expression studies .....	39
3.2.6 Statistical Analysis.....	40
3.3 Results .....	40
3.3.1 Whole-plant physiological responses .....	40
3.3.2 Tissue ionic relations .....	44
3.3.3 Salinity- and hydroxyl radical-induced ion flux kinetics.....	46
3.3.4 Transcriptional changes in the gene expression profile.....	50
3.4 Discussion.....	55
3.4.1 Transcriptional changes in gene expression play a small role in barley adaptive responses to salinity .....	55
3.4.2 Plant tissue tolerance was a dominating component that has determined the overall plant responses to salinity .....	57
3.4.3 Hydroxyl radical-induced K <sup>+</sup> loss is negatively correlated with salinity stress tolerance.....	59
3.5 Conclusion.....	60
Chapter 4    Expressing <i>AtNHX1</i> in barley ( <i>Hordeum vulgare</i> L.) does not improve plant performance under saline condition .....	61
Abstract.....	61
4.1 Introduction .....	62
4.2 Materials and Methods .....	64
4.2.1 Plant Material and Growth Condition.....	64
4.2.2 Cloning of <i>AtNHX1</i> and generating transgenic lines .....	65
4.2.3 Genotyping of <i>35S:AtNHX1</i> in barley .....	65
4.2.4 Determination of transgene expression.....	66
4.2.5 DNA extraction for Southern blot analysis.....	66
4.2.6 Southern blot analysis .....	67
4.2.7 Biomass Measurements .....	67
4.2.8 Na <sup>+</sup> and K <sup>+</sup> content of the leaf.....	67

4.2.9 Gas exchange and water relations .....	68
4.2.10 Statistical analysis .....	68
4.3 Results .....	69
4.3.1 Growth response of barley plants expressing <i>AtNHX1</i> .....	69
4.3.2 Expressing <i>AtNHX1</i> in barley gives no advantage to either enhanced gas exchange or relative water content.....	73
4.3.3. Expressing <i>AtNHX1</i> in barley has not resulted in increased chlorophyll content..	73
4.3.4 Plants expressing <i>AtNHX1</i> showed increased shoot Na <sup>+</sup> and K <sup>+</sup> accumulation .....	73
4.4 Discussion.....	75
4.4.1 Expressing <i>AtNHX1</i> alone in barley has no advantage for biomass .....	75
4.4.2 Barley plants expressing <i>AtNHX1</i> alone has shown no improvement in water relation and gas exchange characteristics .....	77
4.4.3 Sodium and Potassium accumulated in the leaf of barley plants expressing <i>AtNHX1</i> .....	78
Chapter 5 Expressing <i>Arabidopsis thaliana</i> V-ATPase subunit C in barley ( <i>Hordeum vulgare</i> L.) improves plant performance under saline condition by enabling better osmotic adjustment.....	80
Summary.....	80
5.1 Introduction .....	81
5.2 Experimental procedures .....	84
5.2.1 Generating transgenic barley plants expressing <i>VHA-C</i> ( <i>Arabidopsis</i> V-ATPase Subunit C) .....	84
5.2.2 Plant growth conditions .....	85
5.2.3 DNA extraction and PCR analysis.....	85
5.2.4 RNA extraction and gene expression analysis.....	86
5.2.5 Biomass and grain yield.....	86
5.2.6 Pigment content and stomatal conductance .....	87
5.2.7 Na <sup>+</sup> and K <sup>+</sup> content .....	87
5.2.8 Sap osmolality measurements and calculation of contribution of inorganic ions towards leaf osmotic adjustment.....	87
5.2.9 Statistical analysis.....	88
5.3 Results .....	88
5.3.1 Generation of barley plants expressing <i>V-ATPase Subunit C</i> .....	88
5.3.2 Transgenic barley expressing <i>V-ATPase subunit C</i> has improved overall salinity stress tolerance and grain yield .....	90
5.3.3 Transgenic barley expressing <i>V-ATPase subunit C</i> has increased leaf Na <sup>+</sup> and K <sup>+</sup> and these ions contributed towards osmotic adjustment.....	95
5.4 Discussion.....	99
5.4.1 Transgenic <i>VHA-C</i> barley improved growth and grain yield under saline growth condition .....	99

---

5.4.2 Better stomata control contributes to enhanced salinity tolerance in transgenic plants .....	100
5.4.3 Transgenic lines expressing <i>VHA-C</i> have better Na <sup>+</sup> sequestration .....	101
5.4.4 Plants expressing <i>VHA-C</i> showed higher K <sup>+</sup> retention .....	102
5.4.5 Relative contribution of organic and inorganic osmolytes .....	102
Chapter 6    General Discussion .....	104
References .....	113
Appendix .....	134

## List of Figures

**Figure 3.1** Growth responses and agronomical characteristics of three barley cultivars (Numar, Golden Promise, and Naso Nijo) treated with 150 mM NaCl for 4 weeks. A – plant phenotype under control and salt conditions; B – shoot fresh weight; C – shoot dry weight. Open bar - control; closed bar - salt. Mean  $\pm$  SE (n=30).----- 41

**Figure 3.2** Whole-plant physiological characteristics in control- and salt-grown (150 mM NaCl for 4 weeks) plants. A - chlorophyll content (SPAD readings); B - number of necrotic leaves; C – stomatal conductance (Gs); D – relative water content. Open bar - control; closed bar - salt. Mean  $\pm$  SE (n = 10 and 30 for A-C and D, respectively).-----43

**Figure 3.3** Changes in root and shoot sap Na<sup>+</sup> and K<sup>+</sup> content in three barley cultivars contrasting in their salinity stress tolerance during salt stress progression. Plants were grown hydroponically under control conditions until 3 days old, and then exposed to 100mM NaCl treatment. Mean  $\pm$  SE (n=24).----- 45

**Figure 3.4** Kinetics of NaCl- induced net K<sup>+</sup> fluxes measured from the elongation (A) and mature (B) root zones of three barley cultivars in response to 100mM NaCl treatment. Mean  $\pm$  SE (n=6-8). The sign convention is “efflux negative”.----- 47

**Figure 3.5** Kinetics of NaCl- induced net H<sup>+</sup> fluxes measured from the elongation (A) and mature (B) root zones of three barley cultivars in response to 100mM NaCl treatment. Mean  $\pm$  SE (n=6-8). The sign convention is “efflux negative”.----- 48

**Figure 3.6** Hydroxyl radical-induced changes in net K<sup>+</sup> flux measured from the elongation (A) and mature (B) root zones of three barley cultivars. The hydroxyl radical-generated copper ascorbate mix (0.3 mM CuCl<sub>2</sub> and 1 mM Na<sup>+</sup>-Ascorbate) was added at the time indicated by an arrow. Mean  $\pm$  SE (n=6-8). The sign convention is “efflux negative”.----- 49

**Figure 3.7** Expression of barley *NHX1*, *NHX2*, *NHX3*, and *AHA2* transporter genes in leaf and root tissues after 48 h of 100mM NaCl treatment. Mean  $\pm$  SE (n=12-15).----- 51

**Figure 3.8** Expression of barley *RBoHF1* (Respiratory oxidative burst homologue/NADPH oxidase), *RBoHF2*, *SOD2* and *GORK* genes in leaf and root tissues after 48 h of 100mM NaCl treatment. Mean  $\pm$ SE (n=12-15)----- 52

**Figure 4.1** Molecular analysis of *35S:AtNHX1* lines in barley (a) Confirmation of transgene presence by PCR. GP (Golden Promise), L1 (*35S:AtNHX1-1*), L2 (*35S:AtNHX1-2*), N (null line), (-) negative control. (b) Confirmation of transgene expression in transgenic lines by RT-PCR. A multiplex reaction using primers for *AtNHX1* (upper band) and the control gene *Hv-GAP* (lower band) was performed.-----70

**Figure 4.2** Southern blot analysis of two transgenic lines, a null line and Golden Promise. The Genomic DNA was digested with BamHI and HindIII restriction enzymes and probed by hygromycin DNA. The lanes 1-2 and 10-11 correspond to Golden Promise. Lanes 3-4 and 12-13 correspond to null line. Lanes 5-6 and 14-15 correspond to *35S:AtNHX1-2*. Lanes 7- 8 and 16-17 correspond to *35S:AtNHX1-1* with BamHI and HindIII digestion, respectively. Lane 9 and 18 are positive controls.----- 71



**Figure 4.3** Growth and development of three barley *35S:AtNHX1* lines and non-transgenic control cultivar Golden promise (GP) treated with 0 mM, 200 mM NaCl and 250 mM NaCl for five weeks after seedling establishment. *35S:AtNHX1-1* corresponds to line L1, *35S:AtNHX1-2* to L2, and GP to Golden Promise.-----72

**Figure 4.4** Fresh weight (a) and dry weight (b) of two barley *35S:AtNHX1* lines and the null line. Results are mean  $\pm$  SE (n=21-24). Statistical significance designated as  $P \leq 0.05$  (\*),  $P \leq 0.01$  (\*\*) and  $P \leq 0.001$  (\*\*\*)-----72

**Figure 4.5** (a) Stomatal conductance of two barley *35S:AtNHX1* lines and null lines. Results are the mean  $\pm$  SE (n=21-24) (b) Relative water content of two barley *35S:AtNHX1* lines and a null. Results are mean  $\pm$  SE (n=21-24). Statistical significance designated as  $P \leq 0.05$  (\*),  $P \leq 0.01$  (\*\*) and  $P \leq 0.001$  (\*\*\*)-----74

**Figure 4.6** Chlorophyll content (arb. unit) of two *35S:AtNHX1* barley lines and a null line were exposed to 0 mM, 200mM and 250mM NaCl for five weeks after seedling establishment. Results are the mean  $\pm$  SE (n=21-24). Statistical significance designated as  $P \leq 0.05$  (\*),  $P \leq 0.01$  (\*\*) and  $P \leq 0.001$  (\*\*\*)-----74

**Figure 4.7** Leaf  $\text{Na}^+$  (a) and  $\text{K}^+$  (b) content of two *35S:AtNHX1* barley lines and null (used as control) exposed to 200mM and 250mM NaCl for five weeks after seedling establishment. Mean  $\pm$  SE (n=21-24). Mean  $\pm$  SE (n=21-24). Statistical significance designated as  $P \leq 0.05$  (\*),  $P \leq 0.01$  (\*\*) and  $P \leq 0.001$  (\*\*\*)-----75

**Figure 5.1** Genotyping *AtVHA-C* transgenic barley. (a) Detection of transgene presence by PCR using specific primers for *AtVHA-C* and *Hv-VRT2* (internal control); (b) Expression analysis of the *AtVHA-C* gene using RT-PCR with specific primers for the gene and *HvGAP* specific primers (internal control). Lanes *35S:AtVHA-C*-1, 2, 3, & 4 are transgenic *AtVHA-C* barley lines, WT (non-transformed barley cv. WI4330), (-) negative control (water).-----89

**Figure 5.2** Seed size of transgenic lines. The difference in seed size is illustrated for one (OE3 line) of four *35S:AtVHA-C* overexpressing lines under no salt (a, c) and salt stress (b,d) conditions. The other lines show similar results.-----91

**Figure 5.3** Phenotypic difference between non-transformed and overexpressing lines. For simplicity, only two (of four) OE lines are shown. (a-c) Plants treated with 300 mM NaCl for five weeks; (d-f) zero NaCl treatment (no salt stress). The photos were taken 8 weeks after sowing.-----92

**Figure 5.4** Comparative analysis of agronomic characteristics between non-transformed and OE plants. Shoot dry weight (a-c) and plant heights (d-f) are shown for plants grown under zero mM NaCl (no salt; (a, d)) and 300 mM NaCl (saline; b, e) conditions. Panels (c, f) report relative shoot dry weight and plant height, respectively (calculated for salt treated plants as % of control). Results are mean  $\pm$  SE (n=6-15). Measurements are taken 12 weeks after sowing. Asterisks indicate the significant levels of the difference compared with wild type at \* $P \leq 0.05$ ; \*\* $P \leq 0.01$ ; and \*\*\* $P \leq 0.001$ .-----93

**Figure 5.5** Seed number (a-c) and total grain yield (d-f) of OE lines as compared with non-transformed controls. (a, d) – 0 mM NaCl treatment; (b, e) – 300 mM NaCl treatment; (c, f) - relative values (calculated for salt-treated plants as % of control). Results are mean  $\pm$  SE (n=6 to 15). Measurements are taken 12 weeks after sowing. Asterisks indicate the significant levels of the difference compared with wild type at \* $P \leq 0.05$ ; \*\* $P \leq 0.01$ ; and \*\*\* $P \leq 0.001$ .-----94

**Figure 5.6** Difference in the number of chlorotic (a) and necrotic (b) leaves between OE lines and non-transformed barley grown in 300 mM NaCl. Results are mean  $\pm$  SE (n=6 to 15). Measurement was taken 8 weeks after sowing. Asterisks indicate the significant levels of the difference compared with wild type at \*\*\* $P \leq 0.001$ .-----96

**Figure 5.7** Stomatal conductance (a-c) and leaf chlorophyll content (SPAD values; (d-f)) of OE lines as compared with non-transformed barley. (a, d) – 0 mM NaCl treatment; (b, e) – 300 mM NaCl treatment; (c, f) - relative values (calculated for salt-treated plants as % of control). Results are mean  $\pm$  SE (n=6 to 15). Measurement was taken 8 weeks after sowing. Asterisks indicate the significant levels of the difference compared with wild type at \* $P \leq 0.05$ ; \*\* $P \leq 0.01$ ; and \*\*\* $P \leq 0.001$ .-----97

**Figure 5.8** Leaf Na<sup>+</sup> (a-c) and K<sup>+</sup> (d-f) content of OE lines as compared with non-transformed barley. (a, d) – 0 mM NaCl treatment; (b, e) – 300 mM NaCl treatment; (c, f) - relative values (fold changes in salt-treated plants compared with control). Results are mean  $\pm$  SE (n=6 to 15). Measurement was taken 8 weeks after sowing. Asterisks indicate the significant levels of the difference compared with wild type at \* $P \leq 0.05$ ; \*\* $P \leq 0.01$ ; and \*\*\* $P \leq 0.001$ .-----98

## List of Tables

**Table 3.1** The relative gene expression of the gene studied in fold change presented as percentage.-----53

**Table 3.2** The Primers used in the gene expression study and their respective amplicon size.----- 54

**Table 5.1** Seed weight of *AtVHA-C* expressing barley lines and wild type under 0 mM NaCl and 300 mM NaCl growth condition. \*\*\* Significant at  $P \leq 0.001$ . Results are the mean  $\pm$  s.e. (n = 9) ---91

**Table 5.2** Relative contribution of  $\text{Na}^+$ ,  $\text{K}^+$  and  $\text{Cl}^-$  towards total shoot sap osmolality in transgenic barley leaves grown under saline growth condition. Barley lines designated as *35S:AtVHA-C-1* (OE-1), *35S:AtVHA-C-2* (OE-2), *35S:AtVHA-C-3* (OE-3) and non-transformed control (WT). \*Significant compared with WT at  $P \leq 0.01$ .-----99

## **Chapter 1                      General Introduction**

### ***1.1 Salinity's Menace to Agricultural Productivity***

The earth is overwhelmed by water containing high levels of salt, which is estimated to be on average 30 g of NaCl per litre of water. This salt has negatively affected global crop production and continues to challenge productivity, especially within irrigated lands (Flowers 2004). In the bygone times, farmers were aware that part of their lands going out of use as the land is salt-affected; however, the population in these times were small and hence, not concerned to use this lands for agricultural production (Flowers 1999). Almost 6% of the total land cover of the world is affected by high salt content, which is estimated to be more than 800 million hectare of land (FAO 2008). 50% of the irrigated agricultural land, estimated to be 230 million ha is salt affected (Ruan *et al.* 2010). It is estimated that 67% of Australia's agricultural land is exposed to transient salinity (Rengasamy 2006). This may cost the country more than AUD 1330 million per year (Rengasamy 2002). On top of the loss of economic revenue due to salinity stress, it is also important to increase food production by 50%-70% by 2050 in order to feed the ever increasing world population which is projected to reach 9.3 billion (Brown *et al.* 2008; Millar *et al.* 2012; Ruan *et al.* 2010). As it will be difficult to feed this population on the current arable land, it necessitates the expansion of food production to marginal lands, including salt affected land. Due to climate change the mean annual rainfall in subtropical regions is expected to decrease in the coming few decades. This will force farmers to opt for low quality saline water for irrigation (Barrett-Lennard *et al.* 2010). The problem can be further exacerbated by the competition of biofuel crops for productive cultivated land (Valentine *et al.* 2012). Urbanization is additional problem that encroach agricultural land dwindling vast area of arable land. In Australia, agricultural land has declined by 10% in a decade (2001-2009) due to urbanization (Millar *et al.* 2012). In the face of the above mentioned constraints, the option for increasing agricultural production would be either

intensification of agriculture or expansion of arable land where it is required to utilize marginal land. Due to the high cost of input to intensify agricultural production which reduces profitability and sustainability, the use of marginal lands would be of paramount importance. One of the plausible management methods to make saline soils useful for agricultural production is to grow salt tolerant crops. This can be achieved by searching for physiological and genetic traits in plants responsible for tolerance and fortifying these traits into a genotype either using conventional breeding or molecular breeding tools. Developing a salt tolerant crop is a vital measure both to keep the agricultural productivity sustainable and reclaiming the environment.

Salinity can be explained as the excess ion of soluble salt of  $\text{Na}^+$ ,  $\text{Cl}^-$ ,  $\text{Ca}^{+2}$ ,  $\text{SO}_4^{-2}$ ,  $\text{HCO}_3^-$  to the level that it can affect growth and development of plants (Mian 2010). The standard method of measuring salinity is by its electric conductivity (EC). The SI unit for this measuring scale is decisiemens per meter (dS/m). A soil is considered to be saline soil when its EC is greater than 4dS/m (Mian 2010). There are two major types of salinity, namely, primary salinity and secondary salinity. Primary salinity happens when salt is accumulated over a long period of time through natural process such as salt drifted from oceans by wind or water and accumulated on a soil or developed through weathering of rocks. The secondary salinity is anthropogenic (happen by human intervention) like the use of salt rich irrigation water and land clearing (Mian 2010). In sea water, NaCl is the dominant salt and the molar concentration of the major ions in sea water includes  $\text{Na}^+$  approximately 460mM, 50mM for Mg, and 540mM for  $\text{Cl}^-$ . Salinity depends upon the evaporation of a particular area under consideration and/or the precipitation of that given land. If precipitation is high, it will reduce the salt concentration. In desert lands, due to a higher evaporation than precipitation, there is a higher accumulation of salts in the soil. High accumulation of salts also occurs in cultivated lands that get heavy irrigation. This is particularly true where the land is in a dry area that

there will be a high evapo-transpiration as consequence the salt present in the irrigation water accumulates in the soil in due course of time. The repercussion of salinity is a far greater loss of arable land and productivity of most of the important crops as most crops are very sensitive to this environmental stress. Glycophytes is a term used to describe salt sensitive plants and it includes rice (*Oryza sativa*), maize (*Zea mays*), Soybean (*Glycine max*) and beans (*Phaseolus vulgaris*) (Mahajan *et al.* 2005). Although, barley is a glycophyte, it tolerates salinity better than other crops.

## ***1.2 Mechanisms of plant salt tolerance***

### **1.2.1 Osmotic adjustment**

Similar to water stress, salinity stress reduces the ability of the plants to take up water which quickly reduces the growth rate. In a short time after salt stress exposure, the growth rate decreases and the decrease is the same for salt sensitive and salt tolerant species, for example, durum wheat (salt sensitive) vs. bread wheat (relatively salt tolerant) or barley (relatively salt tolerant) vs. triticale cultivars (Munns *et al.* 1995). The temporary drop in the growth rate and rapid recovery to new reduced rate is due to rapid and transient changes in plant water relations (Munns 2002). Rapid and instantaneous change in the growth rate after salinity exposure have been recorded in different crops and similar change have been observed when mannitol, KCl and polyethylene glycol (PEG) are used (Chazen *et al.* 1995; Yeo *et al.* 1991), indicating that it is a non-ion specific response (Munns 2002). This concept was demonstrated by a pressurization technique reported by Passioura *et al.*, (2000) where the plant water status was maintained by an external pressure and the growth rate reduction was not observed after exposure to salinity and the growth rate transiently surged when salt was removed. Similarly the rapid reduction of growth rate in roots as a result of salt stress by affecting the water relations of the root cells was observed (Rodriguez *et al.* 1997). Similar response in roots was observed by using KCl and mannitol (Frensch *et al.* 1994, 1995). Both responses are due to

sudden change in water relations (Munns 2002). In long term of salinity exposure, that is within days' time, whether water status, hormonal regulation or supply of photosynthate is the dominant growth control in saline soil is an issue debated till now. In a pressurization experiment, it was possible to show that in 8 days of salt treatment no effect on growth was found due to the pressure applied (Munns 1993; Termaat *et al.* 1985). This shows that hormonal and other signals are responsible for growth control rather than leaf water deficit or ion toxicity (Munns 2002). Crops produce osmolytes to offset the osmotic pressure imbalance due to salinity stress; the process referred as osmotic adjustment (Hasegawa *et al.* 2000; Munns *et al.* 2015). However, in barley crops, the accumulation of glycine betaine and proline is inversely correlated with potassium efflux (an indicator of salt tolerance) and total amino acid was poorly affected by salt treatment in salt tolerant barley crops (Chen *et al.* 2007a). These authors showed that the accumulation of compatible solutes was more pronounced in salt sensitive barley crops than tolerant cultivars showing that compatible solutes does not have a major role in salt tolerance in barley. Crops like barley or salt includers tend to use inorganic ions as cheap source of osmotic adjustment, as these ions are less energy costly (Munns *et al.* 2015).

### 1.2.2 Sodium exclusion

Exclusion of  $\text{Na}^+$  from the shoot is a crucial mechanism of salinity tolerance in glycophyte plants such as wheat (Colmer *et al.* 2005; Munns *et al.* 2008). Sodium exclusion avoids a high  $\text{Na}^+/\text{K}^+$  ratio in the metabolically active cytosol (Gorham *et al.* 1990), a criteria considered as important for salinity tolerance in plants (Maathuis *et al.* 1999; Shabala *et al.* 2008). Glycophytes have three mechanisms for minimizing  $\text{Na}^+$  accumulation in the shoot: (i) minimizing of entry of  $\text{Na}^+$  into the root; (ii) maximizing  $\text{Na}^+$  efflux from the root; and (iii) restricting  $\text{Na}^+$  transfer to the shoot. Nevertheless, the unidirectional influx of  $\text{Na}^+$  is thermodynamically passive and poorly controlled (Tester *et al.* 2003). Hence, the main

mechanisms for  $\text{Na}^+$  exclusion appears to be active extrusion of  $\text{Na}^+$  back to the rhizosphere and the restriction of  $\text{Na}^+$  loading into the xylem (Cuin *et al.* 2011). In animals, the presence of  $\text{Na}^+$  ATPase facilitates the active transport of  $\text{Na}^+$  out of the cell (Garciadeblas *et al.* 2001). In plants, the active extrusion of  $\text{Na}^+$  from the cytosol is performed by plasma membrane  $\text{Na}^+/\text{H}^+$  antiporter as shown by the lack of  $\text{Na}^+/\text{H}^+$  antiporter activity in the plasma membrane of *sos1* mutant plants (Qiu *et al.* 2002; Qiu *et al.* 2003). Furthermore, sodium extrusion from the cytosol via plasma membrane was also shown by the presence of activity of  $\text{Na}^+/\text{H}^+$  antiporter using highly purified SOS1 protein reconstituted into artificial proteoliposomes, thereby excluding interferences in the  $\text{Na}^+/\text{H}^+$  exchange assay by any other spurious transporter (Quintero *et al.* 2011).

$\text{Na}^+$  extrusion has its own disadvantages. The  $\text{Na}^+$  extruded from the plant build up in the rhizosphere where it creates an increasing gradient. This leads to increased energy cost and as well become thermodynamically more difficult to extrude salt (Cuin *et al.* 2011). This continuous “futile cycling” of  $\text{Na}^+$  could be a problem for the plant and possibly add up to salt toxicity (Tester *et al.* 2003). Under saline growth conditions halophyte plant *Thellungiella halophila*, restricts unidirectional influx of  $\text{Na}^+$  instead of increasing efflux, unlike in *Arabidopsis* (Wang *et al.* 2006). A Similar response was observed in the halophytic monocot *Piccinella tenuiflora* when compared with wheat (Wang *et al.* 2009). If  $\text{Na}^+$  influx into the root is restricted, the energy cost of  $\text{Na}^+$  efflux is reduced. Furthermore, sodium extrusion also results in osmotic stress to plants when salt level builds up in the rhizosphere. The build-up of salt in the reduced apoplastic space- the so called ‘Oertli effect or hypothesis’ could be another possible drawback of a plant’s strategy favouring  $\text{Na}^+$  efflux over reduced influx. The total exclusion of  $\text{Na}^+$  from the plants may not be beneficial in the long term due to the above mentioned reasons (Cuin *et al.* 2011). If  $\text{Na}^+$  is sequestered in the cell vacuole, it could be used as osmoticum to aid cell expansion and also reduces  $\text{Na}^+$  accumulation in the



rhizosphere. To summarize, the water potential in the plant has to be decreased that it could in turn reduce the need to synthesize compatible solutes which is energetically expensive (Cuin *et al.* 2011).

### 1.2.3 Sodium sequestration

The restriction of  $\text{Na}^+$  loading into the xylem does not contribute to salinity tolerance in barley. This has been observed by the tolerant varieties having as high  $\text{Na}^+$  concentration in the xylem as sensitive varieties. The tolerant varieties at the same time maintain high xylem  $\text{K}^+/\text{Na}^+$  ratios and efficiently sequester the excess  $\text{Na}^+$  in the leaves (Shabala *et al.* 2010). Many authors reported that the ability of plants for salinity tolerance is mainly dependent on the exclusion of  $\text{Na}^+$  from the shoot [e.g. (Gorham *et al.* 1990; Munns *et al.* 2008)]. This is possible either by restricting the entry of  $\text{Na}^+$  from the root into the xylem (Davenport *et al.* 2005; Gorham *et al.* 1990), maximizing  $\text{Na}^+$  retrieval from the xylem (Davenport *et al.* 2007), or sending back  $\text{Na}^+$  from the leaf into the phloem (Berthomieu *et al.* 2003). Nevertheless, some species utilize  $\text{Na}^+$  as cheap osmoticum for maintaining cell turgor (Shabala *et al.* 2010), assuming that it can be properly sequestered in the cell vacuole by the tonoplast  $\text{Na}^+/\text{H}^+$  exchanger, NHX (Blumwald 2000). In tolerant barley varieties accumulated  $\text{Na}^+$  is sequestered somewhere in the leaf blade.

The other supporting evidence for this explanation is that it was shown to increase the palisade parenchyma of tolerant barley varieties by NaCl treatment. In addition to keeping the cytosol free of excess  $\text{Na}^+$ , it is also important to compartmentalize  $\text{Na}^+$  for osmotic adjustment in the leaf cells (Shabala *et al.* 2010). Nevertheless, the compartmentalization of  $\text{Na}^+$  in the cell vacuole needs pyramiding of genes involved in the process as well as considering the regulatory components.

#### 1.2.4 Potassium retention

It has been shown that there is a strong negative correlation between  $K^+$  efflux from root and salt tolerance.  $K^+$  efflux from the mature root zone of 3 days old seedling pre-treated with salt is a reliable screening tool for salinity tolerance in barley (Chen *et al.* 2005). Under salt stress, the  $K^+/Na^+$  ratio drops quickly (Maathuis *et al.* 1999). This is due to both accumulation of  $Na^+$  in the cytosol (Zhu 2000) and higher leakage of  $K^+$  from the cell (Shabala 2000; Shabala *et al.* 2003). The  $K^+$  leakage happens because of the NaCl-induced membrane depolarization (Shabala *et al.* 2003). Hence, it's acceptable to use  $K^+/Na^+$  ratio to be used as screening tool for salinity tolerance (Poustini *et al.* 2004; Shannon 1997). However,  $K^+/Na^+$  ratio referred by many breeders is the tissue (root and shoot)  $K^+/Na^+$  ratio rather than the cytosol. Tissue  $K^+/Na^+$  ratio hides the  $Na^+$  compartmentation in the vacuole showing incorrect information which reduces the predictive value of this trait (Chen *et al.* 2005).

The capacity of plant's cell to retain  $K^+$  is important for plant salt tolerance as the ability to exclude or compartmentalize toxic  $Na^+$  (Shabala 2000; Shabala *et al.* 2003). This concept was first tested on seven barley cultivars contrasting in their salt tolerance and proven to be a reliable screening method (Chen *et al.* 2005). In 70 barley cultivars,  $K^+$  flux from the root in response to NaCl treatment was shown to be inversely correlated with relative grain yield, shoot biomass, plant height, net  $CO_2$  assimilation, survival rate and thousand seed weight in glasshouse grown barley plants (Chen *et al.* 2007c). It was shown that the ability of plants to maintain high  $K^+/Na^+$  ratio either by retention of  $K^+$  or by preventing  $Na^+$  from accumulating in leaves is important for salinity tolerance in barley (Chen *et al.* 2007c). For a polygenic trait such as salt tolerance (Flowers 2004), it is difficult to use a single method for 100% predictive value whatever the sensitivity of that method. It was emphasized that a multilateral approach should be used for such complex physiological trait (Chen *et al.* 2007c) such as including other physiological traits with higher correlation with salinity. Using MIFE technique,

measuring both  $K^+$  and  $Na^+$  fluxes would be a highly cost effective and less labour demanding technique to screen barley plants for salt tolerance once the methodological (Cuin *et al.* 2011) and technical issues with  $Na^+$  flux measurement is settled.

### **1.2.5 Reactive oxygen species detoxification**

Reactive oxygen species (ROS) detoxification is also one of the tissue tolerance mechanisms. It was shown that ROS are generated in salinity stressed plant roots and leaves (Miller *et al.* 2008; Mittler 2002). A causal link between ROS and salinity is also becoming evident (Bose *et al.* 2014a). These linkage of salinity with oxidative stress was supported by previous work on  $H_2O_2$  (Maksimovic *et al.* 2013). An increase in ROS production is effected by NADPH oxidase, a cell wall associated peroxidase that produces superoxide by oxidizing NADPH and electron transfer to oxygen (Sagi *et al.* 2006). Hydroxyl radical induced  $K^+$  efflux is mediated by two transport systems, i.e., non-selective cation channel (NSCC) and depolarization-activated  $K^+$  selective outward rectifying channel (Demidchik *et al.* 2003).

## **1.3 Research aims**

The main salt tolerance mechanisms discussed above include osmotolerance, sodium exclusion, sodium sequestration, potassium retention and detoxification of ROS. Among these, the latter three are usually known as components of the tissue tolerance mechanism. The importance and a relative contribution of tissue tolerance mechanisms in the overall salinity stress tolerance is still the matter of debates; the same is true for osmotolerance and  $Na^+$  exclusion. It is also debated of whether adaptation to salinity occurs at transcriptional or post translational level. Some of these gaps in our knowledge were filled in this work. The following major aims addressed were:

- To study the relative contribution of the osmotolerance, sodium sequestration, potassium retention and ROS detoxification in the overall salinity stress tolerance in barley.

- To understand the role of transcriptional and post-translational factors contributing towards above tolerance components.
- To evaluate the feasibility of expression of Arabidopsis  $\text{Na}^+/\text{H}^+$  antiporter *AtNHX1* as a mean for improving salinity tolerance in barley.
- To develop transgenic barley lines expressing *AtVHA-C* (Vacuolar  $\text{H}^+$ ATPase subunit C) to improve biomass and grain yield under saline growth condition.
- To elucidate the prevalent mechanisms of salt tolerance in *AtVHA-C* expressing barley lines.

This chapter has been removed for  
copyright or proprietary reasons.

Chapter 2 Targeting vacuolar sodium sequestration in plant breeding for salinity tolerance (literature review), published work Adem GD., Bose J., Zhou M., and Shabla S. (2015) Targeting vacuolar sodium sequestration in plant breeding for salinity tolerance. In H. Wani and A. Hossain (eds.) Managing salinity tolerance in plants: Molecular and Genomic perspectives (pp. 35-50), Taylor & Francis Catalogue # K23522 ISBN: 978-1-4822-4513-4.

## Chapter 3      Evaluating contribution of ionic, osmotic and oxidative stress components towards salinity tolerance in barley<sup>2</sup>

Getnet Dino Adem<sup>1</sup>, Stuart J Roy<sup>2,3</sup>, Meixue Zhou<sup>1</sup>, John Bowman<sup>1</sup> and Sergey Shabala<sup>1\*</sup>

<sup>1</sup> School of Land and Food, University of Tasmania, Private Bag 54, Hobart Tas 7001, Australia

<sup>2</sup> Australian Centre for Plant Functional Genomics, Private Mail Bag 1, Glen Osmond SA 5064, Australia

<sup>3</sup> University of Adelaide, Private Mail Bag 1, Glen Osmond SA 5064, Australia

### Abstract

Salinity tolerance is a physiologically multi-faceted trait attributed to multiple mechanisms. Three barley (*Hordeum vulgare*) varieties contrasting in their salinity tolerance were used to assess the relative contribution of ionic, osmotic and oxidative stress components towards overall salinity stress tolerance in this species, both at the whole-plant and cellular levels. In addition, transcriptional changes in the gene expression profile were studied for key genes mediating plant ionic and oxidative homeostasis (*NHX*; *RBOH*; *SOD*; *AHA* and *GORK*), to compare a contribution of transcriptional and post-translational factors towards the specific components of salinity tolerance. Our major findings are two-fold. First, plant tissue tolerance was a dominating component that has determined the overall plant responses to salinity, with root K<sup>+</sup> retention ability and reduced sensitivity to stress-induced hydroxyl radical production being the main contributing tolerance mechanisms. Second, it was not possible to infer which cultivars were salinity tolerant based solely on expression profiling of candidate genes at one time point in hydroponics. For the genes studied and the time point selected that

---

<sup>2</sup> This chapter was published in *BMC Plant Biology* 14 (1), 113

transcriptional changes in the expression of these specific genes had a small role for barley's adaptive responses to salinity. For better tissue tolerance, sodium sequestration,  $K^+$  retention and resistance to oxidative stress all appeared to be crucial. Because these traits are highly interrelated, it is suggested that a major progress in crop breeding for salinity tolerance can be achieved only if these complementary traits are targeted at the same time. Also, the importance of post translational modifications calls for salinity research focus being shifted from merely gene expression study to their functional assessment *in planta*.

**Keywords:** stomatal conductance; sodium sequestration; potassium retention; membrane potential; tissue specific responses;  $H^+$ -ATPase; reactive oxygen species; cytosolic ion homeostasis

### **3.1 Background**

The world food supply is endangered by salinity exacerbated by global environmental warming. Agricultural production is already seriously affected by increasing salinity, with estimated economic penalties being in excess of \$12B (Flowers *et al.* 2010; Qadir *et al.* 2008). Creating salt tolerant crop germplasm is, therefore, becoming an urgent imperative (Munns *et al.* 2012; Shabala 2013).

Salinity tolerance in crops is a physiologically multi-faceted trait and is attributed to multiple mechanisms. The key ones include improved osmotic adjustment; minimising  $Na^+$  uptake by roots and/or increasing  $Na^+$  efflux back to the soil; intracellular  $Na^+$  sequestration; potassium retention in the cytosol; tissue-specific  $Na^+$  sequestration; control of xylem ion loading; excluding  $Na^+$  from the shoot; and oxidative stress tolerance (Munns *et al.* 2008; Shabala *et al.* 2012; Zhu 2003). These numerous mechanisms are usually grouped into three major clusters: (i) osmotolerance; (ii) sodium exclusion mechanisms; and (iii) tissue tolerance mechanisms (Munns *et al.* 2008). Despite the significant progress that has been made in elucidating specific details of each of these mechanisms, the relative contribution of the above

components to overall salinity tolerance remain unclear, prompting numerous attempts to overcome the issue by modifying the expression level or function of specific genes by molecular means (Plett *et al.* 2010b; Roy *et al.* 2013).

Crop osmotolerance has long been attributed to a plant's ability to increase *de novo* synthesis of compatible solutes (Hasegawa *et al.* 2000; Shabala *et al.* 2011b). Accordingly, efforts have been made to increase the expression of compatible solute associated genes, most significantly those that catalyse the production of glycine betaine including betaine aldehyde dehydrogenase, encoded by the gene *betB* (Holmstrom *et al.* 2000; Zhou *et al.* 2008), choline dehydrogenase (CDH) encoded by *betA* (Lilius *et al.* 1996) and choline oxidase *codA* (Hayashi *et al.* 1998). Despite these genes being transformed into a large number of species, the success in providing improved crops to the farmers field via this avenue has been very limited (Flowers 2004; Serraj *et al.* 2002).

Another major component of salinity tolerance often targeted in transgenic plants is  $\text{Na}^+$  exclusion by minimising  $\text{Na}^+$  uptake by the root. This component comes to play by the *SOS1* (plasma membrane  $\text{Na}^+/\text{H}^+$  exchanger) along with interacting and phosphorylating proteins *SOS3* and *SOS2* forming a  $\text{Ca}^{2+}$ -dependent signalling cascade (Shi *et al.* 2003; Wu *et al.* 1996). This  $\text{Na}^+$  exclusion process is energised by the plasma membrane  $\text{H}^+$ -ATPase (Palmgren *et al.* 2011). Over-expressing *SOS1*  $\text{Na}^+$  exclusion gene or its homologues has been attempted in several species such as Arabidopsis (Shi *et al.* 2003) or tobacco (Yue *et al.* 2012). However, when  $\text{Na}^+$  is extruded to the medium by this mechanism, it further increases the osmotic and ionic imbalance that in turn causes the stress to be aggravated. Therefore, such a strategy can only be used as an interim solution and cannot, by itself, confer long term salinity stress tolerance under field conditions. Other mechanisms contributing to restricted  $\text{Na}^+$  accumulation in the shoot include reduced  $\text{Na}^+$  loading into the xylem (Munns *et al.* 2008; Tester *et al.* 2003) and increased  $\text{Na}^+$  retrieval from the xylem (Davenport *et al.* 2007), with a



recent study by Munns *et al.*, (2012) showing that the presence of *TmHKT1;5-A* significantly reduced leaf  $\text{Na}^+$  content and increased durum wheat grain yield by 25% compared to near-isogenic lines lacking a *Nax2* locus, expressing this gene. Sodium retrieval from the shoot via its recirculation to the root via phloem is another contributing mechanism (Berthomieu *et al.* 2003), this process is also believed to be mediated by HKT transporters (Garcia-deblas *et al.* 2003). However, it has been argued that excluding  $\text{Na}^+$  from the xylem may not be a plausible mechanism at all times (Shabala *et al.* 2010), as  $\text{Na}^+$  can be used as a cheap osmoticum in species possessing high tissue tolerance mechanisms. Indeed barley over-expressing the *HKT* subfamily 2 gene, *HvHKT2;1*, had higher xylem and leaf  $\text{Na}^+$  content in saline-grown plants and was correlated with increased salt tolerance (Mian *et al.* 2011).

The third component of salinity tolerance is a tissue tolerance, e.g. an ability of plant to safely handle large amounts of sodium accumulated in plant tissues, without any detrimental effects to cell metabolism. Such tissue tolerance can be achieved by efficient  $\text{Na}^+$  sequestration away from the cytosol (where it can affect the metabolic processes of the cell) into vacuole. This is achieved by operation of the tonoplast  $\text{Na}^+/\text{H}^+$  exchanger (Apse *et al.* 1999; Blumwald 2000; Gaxiola *et al.* 1999) that is energised by the vacuolar  $\text{H}^+$ -ATPase (V-ATPase, E.C. 3.6.1.3) and pyrophosphatase (V-PPase, E.C. 3.6.1.1.) (Fukuda *et al.* 2004a). The *Arabidopsis thaliana*  $\text{Na}^+/\text{H}^+$  antiporter gene (*AtNHX1*) was the first plant *NHX* homolog to be cloned (Gaxiola *et al.* 1999). There are six isoforms of *AtNHX* in *Arabidopsis* identified with *AtNHX1* and *AtNHX2* having being highly expressed in many tissues, while *AtNHX3* and *AtNHX4* are exclusively expressed in flowers and roots (Aharon *et al.* 2003; Yokoi *et al.* 2002). Recently, evidence has emerged suggesting that *NHX1* proteins may operate as  $\text{K}^+/\text{H}^+$  exchangers, and that their major role may be not in  $\text{Na}^+$  transport but in regulation of a vacuolar  $\text{K}^+$  content (Barragan *et al.* 2012; Bassil *et al.* 2011b).

Vacuolar Na<sup>+</sup> sequestration is important but not the only mechanism contributing to tissue tolerance. The ability to retain K<sup>+</sup> in cells has recently emerged as an important component of this trait, both in root (Chen *et al.* 2005; Chen *et al.* 2007b; Chen *et al.* 2008; Chen *et al.* 2007d) and leaf (Wu *et al.* 2013) tissues. Strong correlation between K<sup>+</sup> retention ability and plant overall salinity tolerance has been shown in some species (Chen *et al.* 2008; Chen *et al.* 2007d; Smethurst *et al.* 2008) and attributed to the importance of high cytosolic K<sup>+</sup> to suppress activity of caspase-like proteolytic and endonucleolytic enzymes triggering programmed cell death in salt-affected cells (Demidchik *et al.* 2010; Shabala *et al.* 2007). High cytosolic K<sup>+</sup> is also required to maintain metabolic processes such as protein synthesis by enabling tRNA binding to ribosomes (Wyn Jones *et al.* 1979) Also contributing to plant tissue tolerance is reactive oxygen species (ROS) detoxification. It has been shown that significant amounts of ROS are generated in salt-affected plant tissues in both roots and leaves (Miller *et al.* 2008; Mittler 2002), and the causal link between salinity and oxidative stress signalling and reactive species detoxification is becoming evident (Bose *et al.* 2014a). At the same time, attempts to link plants salinity tolerance with the level of antioxidant activity in their tissues appear to be problematic, with reports being highly controversial and ranging from positive to either negative or no correlations with salinity stress tolerance (Bose *et al.* 2014a; Maksimovic *et al.* 2013).

To the best of our knowledge, only one comprehensive attempt to separate the relative contribution of each of above three major components of salt stress was reported in the literature. Using sophisticated whole-plant imaging facilities to determine the area of healthy leaf and the area of senescing leaf in several einkorn wheat accessions, Rajendran *et al.*, (2009) has reported that the most tolerant genotype (judged by relative growth rate under saline conditions) was the *Triticum monococcum* accession, AUS 18755-4 which was not the best performing in any of the above three major mechanisms contributing to salinity tolerance,

namely Na<sup>+</sup> exclusion, osmotic tolerance, and tissue tolerance, compared to the other eleven accessions studied. On the contrary, this variety had an excellent osmotic tolerance (indexed as 0.95 out of 1) and good tissue tolerance but had rather poor ability to exclude Na<sup>+</sup> (indexed as only 0.17). However, given indirect methods of assessing (e.g. whole-plant phenotyping) and the fact that this work has been conducted on salt-sensitive species (wheat), it remains to be answered to what extent these conclusions can be extrapolated to other species.

Barley is one of the most important cereal crops in the world. While being generally classified as relatively salt tolerant (Maas& Grieve 1984), barley germplasm show a great extent of variability in salinity stress tolerance (Chen *et al.* 2007d; Dai *et al.* 2012). We used this opportunity to examine the salt tolerance mechanisms at post-translational level and compared it with changes observed at the transcript level. Unlike to Rajendran *et al.*, (2009), our assay has been conducted not only at the whole plant but also at the cellular level (using the non-invasive microelectrode ion flux measuring (MIFE) technique). We also aimed to compare a contribution of transcriptional and post-translational factors towards the specific components of the overall salinity tolerance. Our results indicate that root K<sup>+</sup> retention ability and increased tolerance to ROS damage were the main contributing tolerance mechanisms. These traits were pronounced at a post translational but not transcriptionally. The overall poor correlation of the change in transcript levels of selected genes in relation to post translational/functional response, demonstrates the importance of post translational modifications *in planta*.

## **3.2 Methods**

### **3.2.1 Plant materials and growth conditions**

#### **3.2.1.1 Glasshouse experiments**

Three barley (*Hordium vulgare* L.) cultivars - Numar (salt tolerant), Naso Nijo (salt sensitive) and Golden Promise (intermediate salinity tolerance) – were used in experiments. Seeds were

obtained from the Tasmanian Institute of Agriculture (TIA) and University of Adelaide Waite Barley Breeders. Seeds were planted into 2L plastic pots containing 70% composted pine bark; 20% coarse sand; 10% sphagnum peat; Limil at 1.8 kg/m<sup>3</sup>, dolomite at 1.8 kg/m<sup>3</sup>). The plant nutrient balance was maintained by adding the slow release Osmocote Plus™ fertilizer (at 6 kg/m<sup>3</sup>), plus ferrous sulphate (at 500 g/m<sup>3</sup>) (Bonales-Alatorre *et al.* 2013a). Two levels of NaCl (0 and 150 mM) were applied in five replications and eight plants were grown in each 2L pot. The plants were grown from seed under controlled greenhouse conditions (temperature between 19 and 26 °C; day length, 12 h; average humidity ~65%) at the University of Tasmania between March 2012 and April 2012. The plants were irrigated with salt free water until seedling establishment (approx. for one week) and then after, the plants were irrigated with the respective salt treatments for four weeks.

### **3.2.1.2 Electrophysiological experiments**

Seeds were surface sterilized by 1.5% (w/v) NaClO and rinsed well with distilled water several times. The seeds were germinated and grown for 3 days in an aerated hydroponic solution containing 0.5 mM KCl and 0.1 mM CaCl<sub>2</sub> in a dark growth chamber at 24 ± 1°C as described elsewhere (Shabala 2000; Shabala *et al.* 2010). Plants were used for measurements when their roots were 60 to 70 mm long.

### **3.2.1.3 Hydroponic experiments**

Barley plants were grown in 1L plastic pot in 25% strength of modified Hoagland solution (Epstein 1972) for 3 days. To salt stress the plants, 100mM of NaCl was then added to the hydroponics solution, and plants were grown for additional 8 days. Root and shoot tissues were sub-sampled on days 1, 2, 4, and 8 after salt application. Control plants were grown in 25% strength of Hoagland solution for the entire duration of experiment.

### 3.2.2 Whole-plant agronomical and physiological characteristics

Shoot fresh (FW) and dry (DW) weight were measured, and relative water content calculated as  $RWC = (FW - DW) / FW$ . Before harvesting, leaf chlorophyll content was measured as a SPAD index using a Minolta Chlorophyll Meter SPAD-502 (Konica Minolta, Osaka, Japan) on the third true leaf, at a position about one quarter of the length of the leaf from the leaf tip. Stomatal conductance ( $G_s$ ) was also measured on the same leaf using a Decagon leaf porometer (Decagon Devices Inc., WA, Australia), under constant light conditions (artificial light of  $150 \mu\text{mol m}^{-2} \text{s}^{-1}$ ). The number of necrotic leaves was also counted at harvest.

### 3.2.3 Tissue sap ion content

$\text{Na}^+$  and  $\text{K}^+$  content in plant tissues was determined by the freeze-thaw method essentially as described elsewhere (Cuin *et al.* 2009). In brief, roots of hydroponically-grown plants were quickly rinsed in 10mM  $\text{CaCl}_2$  to remove apoplastic  $\text{Na}^+$ , blotted dry and then placed into 1.5 mL microfuge tubes and stored at  $-20^\circ\text{C}$ . Shoot samples were harvested at the same time and also frozen at  $-20^\circ\text{C}$ . The frozen samples were thawed and the sap squeezed from the tissue using a pointed glass rod. The sap samples were diluted  $\times 100$  times with distilled water, and  $\text{K}^+$  and  $\text{Na}^+$  content of the sap determined using a flame photometer (MODEL PFP7 Flame photometer, JENWAY, Bibby Scientific Ltd, UK).

### 3.2.4 Non-invasive ion flux measurements

Net fluxes of  $\text{K}^+$  and  $\text{H}^+$  were measured using non-invasive ion measurement technique, MIFE (University of Tasmania, Hobart, Australia) as described in our previous publications (Shabala 2000; Shabala *et al.* 2010). In brief, borosilicate glass microelectrodes with the tip diameter of 2 – 3  $\mu\text{m}$  were pulled, silanised with tributylchlorosilane (Fluka, Catalogue no. 90796), and then filled with appropriate back-filling solution. Electrode tips were then filled with an appropriate Liquid Ion Exchanger (LIX) (Fluka Catalogue no. 60031 for  $\text{K}^+$ ; 95297 for  $\text{H}^+$ ). Microelectrodes were calibrated in a set of pH and  $\text{K}^+$  standards before and after use.

The electrodes were mounted on a 3D-micromanipulator (MMT-5, Narishige, Tokyo, Japan) and the tips of the electrodes were drawn close to each other and positioned 40µm above the root surface. While measuring, the electrodes were moving between two positions (40 and 80 µm) in a 10 s square-wave manner. The CHART software records the potential difference between these two positions and converted it to electrochemical potential difference considering the Nernst slope value obtained during calibration. These potential difference values were converted into ion flux using MIFEFLUX software utilising cylindrical diffusion geometry (see Newman, 2001). Ion fluxes were measured from excised root segments of 3 to 4 day old seedlings from elongation (~ 2 mm from the tip) and mature (~ 10 mm) root zones. Root segments were placed in 10 mL Perspex measuring chamber filled with basic salt medium (BSM; 0.5 KCl mM, 0.1 CaCl<sub>2</sub> mM, pH 5.7 unbuffered) and allowed to equilibrate for ~ 30 minutes. Steady- state ion fluxes were then recorded for 5-10 min, and then the treatment (either 100mM NaCl – for salinity stress; or 0.3mM CuCl<sub>2</sub> + 1mM ascorbate – for ROS stress) was administered.

### 3.2.5 Gene Expression studies

Barley cultivars were grown hydroponically until the third leaf was fully emerged (plants were approximately 15 days). The growth solution was changed every seven days. Plants were treated with 100mM NaCl for 48 hours, their roots and leaves harvested, and snap frozen with liquid nitrogen. Total RNA was extracted following the method of Chomczynski, (1993), using TRIzol reagent (Invitrogen, Carlsbad, CA, USA). Ambion's DNA-free (Madison, WI, USA) was used to remove contamination of genomic DNA. To synthesise cDNA, Invitrogen's superscript III Reverse Transcriptase kit with an oligo(dT)<sub>20</sub> primer was used, following the manufacturer's instructions. Quantitative Real-time PCR was performed as described in (Burton *et al.* 2008) using a RG6000 Rotor-Gene real time thermal cycler (Corbett Research, Sydney) and SYBR<sup>®</sup> green PCR reagent (Bio-Rad Laboratories,

Gladesville). Primers were designed to determine the expression of a number of key genes involved in  $\text{Na}^+$  compartmentation and ROS detoxification and included members of the *NHX*, *AHA*, *RBoHF*, *SOD* and *GORK* family of genes. Primer sequences can be found in Table 3.2. Normalization of the test gene transcript was relative to the control gene (*GAPdH2*).


### **3.2.6 Statistical Analysis**

All the values in this manuscript are presented as mean value  $\pm$  SE. For mean comparison and statistical significant level paired wise t-test in all possible combination of the treatments was employed using SPSS software version 20 (IBM support portal, USA).

## **3.3 Results**

### **3.3.1 Whole-plant physiological responses**

Salinity stress significantly affected plant growth and biomass production (Fig. 3.1A), resulting in a three- and five-fold reduction in the fresh weight in the tolerant variety Numar and the sensitive cultivar Naso Nijo, respectively (FW; Fig. 3.1B). The cultivar Golden Promise displayed an intermediate salinity tolerance with a 4-fold FW reduction in biomass production under salt stress. All the differences were significant at  $P < 0.001$ .




**Figure 3.1** Growth responses and agronomical characteristics of three barley cultivars (Numar, Golden Promise, and Naso Nijo) treated with 150 mM NaCl for 4 weeks. A – plant phenotype under control and salt conditions; B – shoot fresh weight; C – shoot dry weight. Open bar - control; closed bar - salt. Mean  $\pm$  SE (n=30).

Four weeks of salinity stress also resulted in a reductions in leaf chlorophyll content (Fig. 3.2A). This reduction in chlorophyll varied significantly among cultivars, with the salt



sensitive Naso Nijo exhibiting the highest reduction in chlorophyll content (a 4-fold reduction from  $29 \pm 0.7$  to  $7.9 \pm 1.6$  arbitrary units; significant at  $P < 0.001$ ). The salt-tolerant cultivar Numar, however, increased its leaf chlorophyll content by  $\sim 10\%$  (significant at  $P < 0.05$ ), while the intermediate salt tolerant Golden Promise showed no significant reduction in chlorophyll content. The chlorophyll content of the salt grown Numar leaves was 4-fold greater than that measured in salt stressed Naso Nijo leaves (all the differences are significant at  $P < 0.01$ ; Fig. 3.2A). The salt-sensitive Naso Nijo had twice as much necrotic leaves as Numar and Golden Promise (Fig 3.2B)


Salinity stress also resulted in a significant ( $P < 0.01$ ) decline in stomatal conductance ( $G_s$ ; Fig 3.2C) and shoot water content (RWC; Fig. 3.2D). The highest decline in  $G_s$  was measured in the sensitive cultivar Naso Nijo (a five-fold reduction from  $53.2 \pm 4.7$  to  $10.4 \pm 2.3$   $\text{mmol m}^{-2} \text{s}^{-1}$ ). Surprisingly, the intermediate cultivar Golden Promise outperformed salt tolerant variety Numar (2-fold vs 4.5-fold  $G_s$  reduction, respectively; both significant at  $P < 0.01$ ). However, due to initially higher  $G_s$  values in control conditions for Numar, the difference in  $G_s$  between salt-grown Golden Promise and Numar genotypes was not statistically significant (at  $P < 0.05$ ). As a result, both these varieties retained significantly ( $P < 0.01$ ) more water in the shoot compared with salt-sensitive Naso Nijo (Fig. 3.2D).



**Figure 3.2** Whole-plant physiological characteristics in control- and salt-grown (150 mM NaCl for 4 weeks) plants. A - chlorophyll content (SPAD readings); B - number of necrotic leaves; C – stomatal conductance (Gs); D – relative water content. Open bar - control; closed bar - salt. Mean  $\pm$  SE (n = 10 and 30 for A-C and D, respectively).

### 3.3.2 Tissue ionic relations

Hydroponic experiments demonstrated that salt-tolerant Numar accumulated less  $\text{Na}^+$  in the root (Fig 3.3A) compared with two other cultivars, throughout the whole salt stress period. At the same time, Numar plants had twice as much  $\text{Na}^+$  in the shoot sap compared with other cultivar on day 4 (Fig. 3.3B), however, similar shoot  $\text{Na}^+$  were observed between all three cultivars by day 8 (Fig. 3.3B). There was no significant difference in either root or shoot  $\text{Na}^+$  accumulation between the three cultivars when grown in the absence of salt (data not shown). Salinity stress resulted in a rapid and progressive decline in root  $\text{K}^+$  content in salt-sensitive cultivar Naso Nijo (a 2-fold decrease from  $128.4 \pm 31$  to  $62.4 \pm 2.7$  mM over 8 days of 100 mM NaCl treatment; Fig 3.3C), while over the same time period, root  $\text{K}^+$  content did not change significantly in salt-tolerant cultivar Numar. The root  $\text{K}^+$  content in the intermediate salt tolerant Golden Promise also declined substantially, but not to the same extent as in Naso Nijo (Fig 3.3C). Interestingly, while root  $\text{K}^+$  content dropped sharply in cultivars Naso Nijo and Golden promise, their shoot  $\text{K}^+$  content increased approximately 2 fold over the same time period (Fig. 3.3D), suggesting a possible retranslocation of  $\text{K}^+$  from root to shoot. This observation was not seen for Numar (Fig 3.3D). The mean values for root  $\text{K}^+$  content in controls were  $111.2 \pm 11.6$ ,  $153 \pm 4.5$  and  $146.7 \pm 1.9$  mM, for Naso Nijo, Golden Promise and Numar, respectively.




**Figure 3.3** Changes in root and shoot sap Na<sup>+</sup> and K<sup>+</sup> content in three barley cultivars contrasting in their salinity stress tolerance during salt stress progression. Plants were grown hydroponically under control conditions until 3 days old, and then exposed to 100mM NaCl treatment. Mean  $\pm$  SE (n=24).

### 3.3.3 Salinity- and hydroxyl radical-induced ion flux kinetics


Similar to our previous reports on barley (Chen *et al.* 2005; Chen *et al.* 2007d; Maksimovic *et al.* 2013), acute NaCl treatment induced massive  $K^+$  efflux from the epidermal cells of the root (Fig. 3.4). Epidermal cells in elongation zone had approximately 5-fold higher  $K^+$  efflux when compared with those in mature zone (Figs 3.4A and B, respectively; significant at  $P < 0.01$ ). Responses of cultivar Golden Promise and Numar were very similar (no difference at  $P < 0.05$ ) while the magnitude of NaCl-induced  $K^+$  leak from the roots of the salt-sensitive Naso Nijo was approximately 4-fold greater than the response seen in the other two cultivars (Fig 3.4; significant at  $P < 0.05$ ).

Salinity treatment has also induced a rapid (within one minute) net  $H^+$  efflux from both the elongation and mature root zones (Fig. 3.5). This activation was strongest for Golden Promise, followed by Numar and then Naso Nijo (Fig. 3.5). The difference in  $H^+$  efflux amongst cultivars was significant at  $p < 0.01$ ; and so was the difference between elongation and mature zones (Fig. 3.5A & B). The NaCl-induced  $H^+$  efflux was more pronounced in the elongation zone in comparison to the mature zone.


Consistent with previous observations (Cuin *et al.* 2007; Demidchik *et al.* 2010), addition of hydroxyl radical-generating Cu/ascorbate mix has also triggered massive  $K^+$  efflux from plant roots (Fig. 3.6). Similar to NaCl stress, responses from elongation zone for each cultivar was significantly ( $P < 0.05$ ) stronger compared with mature root zone (Fig. 3.6A and 6B, respectively). ROS induced  $K^+$  efflux was higher in the salinity sensitive cultivar Naso Nijo followed by intermediate Golden Promise variety and then the tolerant cultivar Numar (Fig 3.6B; significant at  $P < 0.05$ ).



**Figure 3.4** Kinetics of NaCl- induced net K<sup>+</sup> fluxes measured from the elongation (A) and mature (B) root zones of three barley cultivars in response to 100mM NaCl treatment. Mean  $\pm$  SE (n=6-8). The sign convention is “efflux negative”.



**Figure 3.5** Kinetics of NaCl- induced net H<sup>+</sup> fluxes measured from the elongation (A) and mature (B) root zones of three barley cultivars in response to 100mM NaCl treatment. Mean  $\pm$  SE (n=6-8). The sign convention is “efflux negative”.



**Figure 3.6** Hydroxyl radical-induced changes in net K<sup>+</sup> flux measured from the elongation (A) and mature (B) root zones of three barley cultivars. The hydroxyl radical-generated copper ascorbate mix (0.3 mM CuCl<sub>2</sub> and 1 mM Na<sup>+</sup>-Ascorbate) was added at the time indicated by an arrow. Mean  $\pm$  SE (n=6-8). The sign convention is “efflux negative”.



### 3.3.4 Transcriptional changes in the gene expression profile

The expression of a number of key genes involved in  $\text{Na}^+$  compartmentation and ROS related were examined in hydroponically-grown plants (Figs 3.7, 3.8; Table 3.1). The expression pattern of *GORK* was also examined, given the essential role cytosolic  $\text{K}^+$  retention plays in salinity tolerance in barley (Chen *et al.* 2005; Chen *et al.* 2007d) .with the GORK channel having a key role in this process (Shabala *et al.* 2008).


As shown in Fig. 3.7 and 3.8, there was a large range in the transcriptional response in the eight genes studied to salt application in both leaves and roots. However no clear patterns of gene expression could be observed which could explain the difference in tolerance between the three cultivars. Of three homologues to the *AtNHX* gene, which was originally suggested to encode a protein involved in  $\text{Na}^+$  sequestration into the vacuoles, a 2-fold increase in *NHX1* transcript level was measured both in the leaf and root of the salt sensitive NasoNijo (Fig 3.7). No significant changes in *NHX1* expression were measured in the leaf and root of the salt-tolerant Numar plants, while in the intermediate Golden Promise significant *NHX1* upregulation was only observed in the leaves (Fig 3.7). *NHX2* transcript levels went down in both leaves and roots in cultivars Numar and Golden Promise but not in salt-sensitive NasoNijo (Fig. 3.7), while no clear patterns were observed for *NHX3* transcripts.

During salinity treatment there was a relative increase in the expression of the barley homologue *AHA2* in the roots of both Numar and Naso Nijo, while a significant induction was observed only in the leaves of Naso Nijo (Fig. 3.7). No significant ( $P < 0.05$ ) changes in *AHA2* transcript levels were detected in Golden Promise plants.


The relative gene expression of barley homologue of *RBoHF1*, important in encoding for NADPH oxidase was down-regulated significantly ( $P < 0.01$ ) by about 2-fold in the leaves of all the cultivars after salt application (Fig. 3.8), however, no clear patterns in the roots were detected. *RBoHF2*, also involved in the production of NADPH oxidase was only slightly

down-regulated after salt application (significant for Numar in leaves and Naso Nijo and Golden Promise in roots; Fig 3.8). The barley homologue of *SOD2* relative gene expression showed small increase in leaves of Golden Promise cultivar but no significant changes for any other varieties in either tissue (Fig. 3.8).

The largest and most striking difference was observed for NaCl-induced changes in the relative transcript abundance of the barley homologue of *GORK* in barley roots, with 4 to 5-fold increase reported for all cultivars (Fig. 3.8; all significant at  $P < 0.01$ ). In leaves, small but significant (at  $P < 0.05$ ) increase in *GORK* transcript level was measured in the salt-sensitive Naso Nijo while the *GORK* transcripts in the leaf of salt-tolerant Numar were reduced (Fig. 3.8).



**Figure 3.7** Expression of barley *NHX1*, *NHX2*, *NHX3*, and *AHA2* transporter genes in leaf and root tissues after 48 h of 100mM NaCl treatment. Mean  $\pm$  SE (n=12-15).



**Figure 3.8** Expression of barley *RBoHF1* (Respiratory oxidative burst homologue/NADPH oxidase), *RBoHF2*, *SOD2* and *GORK* genes in leaf and root tissues after 48 h of 100mM NaCl treatment. Mean  $\pm$ SE (n=12-15)

**Table 3.1** The relative gene expression of the gene studied in fold change presented as percentage.

Gene Name	% Fold Change (p-value)			% Fold Change (p-value)		
	Leaf			Root		
	Cultivar			Cultivar		
	NN	GP	Numar	NN	GP	Numar
<i>Hv-NHX1</i>	200	160	-95	171	-89	178
	(0.005)	(0.02)	(NS)	(0.01)	(NS)	(NS)
<i>Hv-NHX2</i>	-94	-80	-68	141	-64	-76
	(NS)	(0.0001)	(0.0001)	(0.03)	(0.0001)	(NS)
<i>Hv-NHX3</i>	104	118	-73	-76	-58	137
	(NS)	(0.01)	(0.0009)	(NS)	(0.0001)	(NS)
<i>Hv-RBoHF1</i>	-60	-39	-49	-81	-57	143
	(0.0001)	(0.0001)	(0.0001)	(NS)	(0.0001)	(NS)
<i>Hv-RBoHF2</i>	131	-78	-74	-76	-58	134
	(NS)	(NS)	(0.0001)	(0.05)	(0.0001)	(NS)
<i>Hv-PMHATPase</i>	212	-84	99	161	117	169
	(0.0001)	(NS)	(NS)	(0.0005)	(NS)	(0.02)
<i>Hv-SOD2</i>	107	136	97	-92	-83	113
	(NS)	(0.0001)	(NS)	(NS)	(0.05)	(NS)
<i>Hv-GORK</i>	137	129	-74	339	375	487
	(0.006)	(NS)	(0.0001)	(0.0001)	(0.0001)	(0.0001)
NS-Non-significant; - sign indicate down-regulation						

**Table 3.2** The Primers used in the gene expression study and their respective amplicon size.

Gene Name	Forward Primer	Reverse Primer	Amplicon Size (bp)
<i>Hv-NHX1</i>	TGCATATCTACCA- GTGCTTAT	GGTTCAAGACACA- AGTTCAGT	184
<i>Hv-NHX2</i>	GGTTTTTCGGCTTG- CTGACTAA	CATTGGGCGCATG- AACTTATC	238
<i>Hv-NHX3</i>	TGAGCCGAACATT- ACTGTGAT	ACGAGCTTACCTT- TCAATACA	127
<i>Hv-RBoHF1</i>	TTACAACATGGAC- CTGCGTCCCTACA	TGCCTTGGTCAGA- CACTCAGCTGCAT	206
<i>Hv-RBoHF2</i>	TATGCGGAGTCCC- GCAGAAAGATG	TGTACTGTACTCC- CCCTGCCTGTGT	212
<i>Hv-PMHATPase</i>	CTTGGTTATCGCC- TTCCTTC	GCCACTCAGCACA- AATCG	184
<i>Hv-SOD2</i>	CTTGAAGGACACC- GACTTGC	CTCAAAAAGCCA- AATGACAGTG	140
<i>Hv-GORK</i>	CCACACGAGGCGA- AGAAG	GAGGAATCCACAG- CATCACC	194
<i>Hv-GAPdH2</i>	GTGAGGCTGGTGC- TGATTACG	TGGTGCAGCTAGC- ATTTGAGAC	Control gene

### **3.4 Discussion**

#### **3.4.1 Transcriptional changes in gene expression play a small role in barley adaptive responses to salinity**

Vacuolar sodium sequestration is essential to avoid  $\text{Na}^+$  cytotoxicity, and increased activity of *NHX* tonoplast  $\text{Na}^+/\text{H}^+$  exchangers was shown to enhance salinity tolerance in plants (Apse *et al.* 1999; Blumwald *et al.* 2000). However, in the work presented here a 2-fold increase in *NHX1* transcript level was measured in both leaves and roots of salt-sensitive Naso Nijo (Fig 3.7), while no significant (at  $P < 0.05$ ) changes were measured in salt-tolerant Numar plants in either tissue. Consistent with this, *NHX2* transcript levels went down in both tissues in cultivars Numar and Golden Promise but not in salt-sensitive Naso Nijo (Fig. 3.7). Thus, it appears that changes in transcript levels of the *NHXs* have a little role (and low predictive value) on the salinity stress tolerance in barley in our experiments and changes in expression cannot be used to predict salt tolerance.

Several possible explanations should be considered for these observations. First, as described above, there is recent evidence that the major role of *AtNHX1* may be in  $\text{K}^+$  but not  $\text{Na}^+$  transport (Bassil *et al.* 2011b), thus the genes studied here may not encode proteins involved in  $\text{Na}^+$  compartmentation. Second, the higher NaCl induced expression of *NHXs* in the shoots of the salt-sensitive Naso Nijo could be to compensate for its inability to prevent  $\text{Na}^+$  delivery to the shoot (e.g. a need to deal with the consequences of cytosolic  $\text{Na}^+$  accumulation). However, this explanation is unlikely as the shoot  $\text{Na}^+$  content of the salt tolerant Numar was higher than that of NasoNijo between days 2 and 4 of salt stress (the time when gene expression of the *NHXs* were assessed). Another explanation is that post-translational modifications such as protein folding are much more important than gene expression to have a proper functional response – the proteins are already present at desired concentrations (therefore gene expression is not required), however, need to be activated by

post-translational modifications, a process which allows a plant to respond faster to stress than relying solely on gene expression. It should also be noted that only one time point was used for the expression studies, and expression of genes encoding salt tolerance gene has been shown to fluctuate between days (e.g. *HVP10* in (Shavrukov *et al.* 2013)). Lastly, it should also be noted the *NHX1* and *NHX2* genes in barley are homologues of *AtNHX1* and, while having similar nucleotide sequence may not necessarily confer the protein(s) function as *AtNHX1* in Arabidopsis.

The above notion that the changes in transcript levels do not always confer plant salinity stress tolerance is further corroborated by study of plant oxidative stress responses. No significant difference in the relative expressions of *RBoHF1* and *RBoHF2* which encode the NADPH oxidase/*Nox* (one of the major sources of ROS production under saline conditions; (Bolwell *et al.* 1997; Kawano 2003)) were found in roots of the salt sensitive NasoNijo and salt tolerant Numar cultivars (Fig. 3.8). Similarly, salt stress did not result in any significant changes in *SOD* expression level in either root or leaf tissues of these varieties (Fig. 3.8). At the same time, NasoNijo roots were twice more sensitive to ROS treatment (Fig. 3.6B).

Other evidence comes from comparing the NaCl-induced net  $H^+$  fluxes measured in barley roots (Fig 3.5) with the changes in the *AHA* (encoding plasma membrane  $H^+$ -ATPase) transcript levels (Fig 3.7).  $H^+$ -ATPase activity is indispensable for maintaining membrane potential (Palmgren *et al.*, (2011), and intrinsically higher  $H^+$ -ATPase activity was shown to correlate with salinity tolerance in barley (Chen *et al.* 2007d). In this study, the Golden Promise cultivar showed consistently higher proton efflux in both elongation and mature zone compared with two other cultivars (Fig. 3.5A, 3.5B). However, the expression of the *AHA* barley homologue was shown to be lower compared to the other two cultivars suggesting that posttranslational modifications may result in higher  $H^+$  pumping activity in Golden Promise cultivar.

### **3.4.2 Plant tissue tolerance was a dominating component that has determined the overall plant responses to salinity**

The tolerant cultivar Numar maintained root  $K^+$  content at a constant level throughout the eight day salt stress period, while the two other varieties showed a progressive decline in root  $K^+$  content (with the greatest decline in the salt-sensitive NasoNijo) (Fig. 3.3C). This was further corroborated in MIFE experiments measuring NaCl-induced  $K^+$  efflux from barley roots (Fig. 3.4). The rationale behind this experiment is that when  $Na^+$  is absorbed from extracellular space, the membrane gets depolarised and this depolarisation of cell membrane initiates potassium leak as a result of the activation of depolarisation activated outward-rectified potassium channels (KOR) (Chen *et al.* 2007d; Shabala *et al.* 2006). This reduces the  $K^+$  content in the cytosol negatively affecting cell metabolism (Shabala *et al.* 2008) and in turn brings about programmed cell death (Demidchik *et al.* 2010; Shabala 2009). Potassium loss from the epidermal cell in the elongation and mature zone of three barley plants was studied and the highest potassium loss was observed in the salt sensitive cultivar NasoNijo followed by the tolerant cultivars (Numar and GP) in both zones (Fig. 3.4 A, B), reflecting overall tolerance estimated by agronomical (biomass accumulation; Fig. 3.1) and whole-plant physiological (chlorophyll content; Fig. 3.2) characteristics. At the same time, osmotolerance appears to be not central to the overall plant performance under saline conditions. Indeed, the intermediate salt tolerant Golden Promise outperformed the salt-tolerant cultivar Numar, having highest  $G_s$  and shoot water content values under saline conditions (Fig. 3.2). This corroborates the point made by Rajendran *et al.*, (2009) that superiority in one salt tolerant component does not guarantee an enhanced overall salt tolerance performance.

Gas exchange ( $G_s$ ) could be considered as a yield determinant and a valuable tool as a physiological trait that can readily be used as a breeding tool (Lu *et al.* 1998; Richards 2000). Carbon entry and transpiration occurs using open stomata which both helps to increase



photosynthesis and nutrient absorption from the growth media. However, under saline conditions, the stomata tend to close to prevent water loss as it is a scarce resource due to osmotic imbalance. Thus, higher  $G_s$  values will ensure better  $CO_2$  assimilation ability (and, hence, higher yield) only when plants have a biochemical and physiological capacity to fully utilise it. This means that higher  $G_s$  values must be complemented by higher shoot tissue tolerance, to ensure efficient leaf photochemistry under saline conditions.

To our great surprise, the expression of the barley homologue of *GORK* gene showed high up-regulation in all cultivars. This is counterintuitive, as the opposite effect would be hypothesised due to the very strong correlation between barley salinity stress tolerance and its ability to prevent NaCl-induced  $K^+$  leak from roots (Chen *et al.* 2005; Chen *et al.* 2007b; Chen *et al.* 2007d). These findings, however, are fully consistent with the above notion that transcriptional changes appear to be causally unrelated to functional plant responses. It could be suggested that the post-translational phenomena of assembly might play a role in the regulation of this gene as it is fundamental for the pore formation and electric activity to form tetramer of its  $\alpha$ -subunit (Dreyer *et al.* 1997). Furthermore, the availability of the  $\alpha$ -subunits and its functional tetrameric assembly is of paramount importance for the cell (Dreyer *et al.* 2004).

Another fact supporting the essential role of tissue tolerance in overall plant performance under saline conditions could be found in kinetics of  $Na^+$  accumulation in the shoot (Fig. 3.3). Consistent with previous reports (Bose *et al.* 2014a; Mian *et al.* 2011; Shabala *et al.* 2010), the tolerant cultivar Numar tends to send more  $Na^+$  to the shoot via the transpiration stream and use it as a cheap osmoticum to maintain shoot turgor, while sensitive variety Naso Nijo delayed this process favouring  $Na^+$  accumulation in the roots (Fig. 3.3A, B). Despite this, Numar plants were capable to maintain higher chlorophyll levels while massive chlorosis was observed in Naso Nijo variety (Fig 3.3). As the volume of the leaf epidermis is small; the

explanation for  $\text{Na}^+$  sequestration would be vacuolar compartmentalisation in the mesophyll cells. Consistent with this, salinity treatment caused an increase in the size of palisade parenchyma cells; this increase was much higher in the tolerant barley cultivars (Shabala *et al.* 2010). A similar phenomena is widespread in halophytes (Shabala 2013), species considered to be most tolerant to salt stress. Halophytes also use additional tissue tolerance mechanism such as salt bladders to accumulate excess  $\text{Na}^+$  away from the photosynthetic tissue.

### **3.4.3 Hydroxyl radical-induced $\text{K}^+$ loss is negatively correlated with salinity stress tolerance**

In addition to the two above components of the tissue tolerance mechanism, namely better  $\text{K}^+$  retention by roots and more efficient vacuolar  $\text{Na}^+$  sequestration in shoots, salinity stress tolerance in barley is also correlated strongly with its ability to prevent hydroxyl radical - induced  $\text{K}^+$  loss (Fig 3.6). The highest potassium efflux was observed from the salt sensitive cultivar Naso Nijo, followed by Golden Promise (intermediate) and Numar (tolerant). The hydroxyl radical-induced effects were higher in the mature root zone (Fig. 3.6A & B). These findings are consistent with previous reports for root sensitivity to  $\text{H}_2\text{O}_2$  (Chen *et al.* 2007d; Maksimovic *et al.* 2013) suggesting that sensitivity to oxidative stress is an essential component of the tissue tolerance mechanism.

An increase in ROS production under salinity stress is attributable to activity of NADPH oxidase, a cell wall-associated peroxidase that generates  $\text{O}_2^{\cdot-}$  by oxidizing NADPH and transferring the electron to oxygen ( $\text{O}_2$ ) (Sagi *et al.* 2006). Accordingly, the regulation of the gene at the transcript level was studied. The relative gene expression at the transcript level of the barley homologue *RBoHF* gene for all of the cultivars was significantly down-regulated but only in leaves but not roots (Fig. 3.8). This down-regulation was least in salt-sensitive NasoNijo, suggesting a possible causal link between NADPH oxidase activity and tissue tolerance mechanisms in salinised mesophyll tissues. However, unlike the leaf, in the root the

transcript abundance was non-significant, except for Golden Promise (Fig. 3.8). The plausible explanation for this could be the post-translational modification such as activation by  $\text{Ca}^{2+}$  owing to the EF-hand motif for binding  $\text{Ca}^{2+}$  (Sagi *et al.* 2006).

Potassium efflux induced by hydroxyl radicals is mediated by two major transport systems: (i) non-selective cation channel (NSCC) and (ii) depolarization-activated  $\text{K}^+$ -selective outward rectifying channel (Demidchik *et al.* 2010; Demidchik *et al.* 2003). High cytosolic  $\text{K}^+$  levels are also essential to suppress activity of caspase-like proteases and endonucleases, both in mammalian (Hughes *et al.* 1998, 1999) and plant (Demidchik *et al.* 2010; Shabala *et al.* 2007) systems. Decrease in the cytosolic  $\text{K}^+$  pool in plant roots may result in activation of these catabolic enzymes triggering programmed cell death (PCD), especially in sensitive root apex cells (Fig. 3.6).

### **3.5 Conclusion**

In this study, it was possible to show that much of the genes under scrutiny, at the transcriptional level, has lower explanatory value towards the contribution of this gene towards plants adaptive responses to salinity. At the post-translational (e.g. functional) level, tissue tolerance was the dominant mechanism of the salt tolerance. For better tissue tolerance, sodium sequestration,  $\text{K}^+$  retention and resistance to oxidative stress all appeared to be crucial. These traits seem to be highly interrelated, as cytosolic  $\text{K}^+$  retention is essential for the optimal vacuolar  $\text{H}^+$  pump operations (Shabala 2013) required to fuel  $\text{NHX Na}^+/\text{H}^+$  exchanger to enable  $\text{Na}^+$  sequestration. Cytosolic  $\text{K}^+$  homeostasis, in its turn, is strongly affected by the sensitivity of plasma membrane transporters to ROS and the plant's ability to prevent stress-induced hydroxyl radical accumulation in stressed tissues. Thus, it appears that a major progress in crop breeding for salinity tolerance can be achieved only if these complementary traits are targeted at the same time.

## Chapter 4      Expressing *AtNHX1* in barley (*Hordeum vulgare* L.) does not improve plant performance under saline condition<sup>3</sup>

Getnet D. Adem<sup>1</sup>, Stuart J. Roy,<sup>2,3</sup> Darren C. Plett<sup>2,3</sup>, Meixue Zhou<sup>1</sup>, John P. Bowman<sup>1</sup> and, Sergey Shabala.<sup>1</sup>

<sup>1</sup>School of Land and Food, University of Tasmania, Private Bag 54, Hobart, Tas 7001, Australia

<sup>2</sup> Australian Centre for Plant Functional Genomics, Private Mail Bag 1, Glen Osmond, SA 5064, Australia

<sup>3</sup> School of Agriculture, Food and Wine, University of Adelaide, Private Mail Bag 1, Glen Osmond, SA 5064, Australia

Correspondence: Sergey Shabala, address : School of Land and Food, University of Tasmania, Private Bag 54, Hobart, Tas 7001, Australia (Tel. +61 3 6226 7539; Fax +61 3 6226 7444; email Sergey.Shabala@utas.edu.au)

### Abstract

Salinity is one of the main causes of yield loss in agricultural crop production worldwide. Numerous attempts have been made to produce salinity tolerant crops by manipulating the expression of genes involved in ion transport; yet, the reported results are often rather controversial. In this work, Arabidopsis *AtNHX1* gene encoding vacuolar Na<sup>+</sup>/H<sup>+</sup> exchanger was expressed in barley (*Hordeum vulgare* L. cv. Golden Promise) plants grown under saline conditions. Transgenic plants were compared to null segregants for biomass, water content, gas exchange, and Na<sup>+</sup> and K<sup>+</sup> content of the leaf. It was shown that most of the lines expressing *AtNHX1* have no significant difference to null segregants, suggesting the lack of beneficial effect of expressing the *AtNHX1* gene in barley. The lack of phenotype might be

---

<sup>3</sup> This paper was published in *Plant Growth Regul.* 1-9.

explained by: (i) low level of activity of vacuolar  $H^+$ -inorganic pyrophosphatase and vacuolar  $H^+$ -ATPase resulting in an insufficient proton gradient required for vacuolar  $Na^+/H^+$  exchanger; (ii) the inability of transgenic plants to prevent a passive leak of sodium via  $Na^+$  permeable slow activating (SV) and fast activating (FV) channels; (iii) insufficient ATP pool to support  $H^+$  pumping activity; and (iv) the fact that the *AtNHX1* protein might not been properly folded, inactive or incorrectly targeted. Overall, this study suggests that, while targeting vacuolar  $Na^+$  sequestration, at least several genes need to be pyramided to ensure the efficiency of the above process. It also suggests the need to verify the proper folding and targeting of the protein in transgenic plants.

**Keywords:** barley, *AtNHX1*, Salinity, vacuolar sequestration, vacuolar channels, proton pumping

## 4.1 Introduction

Saline soils cover 950 million ha, accounting for 10% of the land surface. It has been estimated that 50% (230 million ha) of irrigated land are affected by salt (Ruan *et al.* 2010) and is responsible for an estimated loss of \$US27 billion per annum (Qadir *et al.* 2014). In Australia, 67% of the agricultural land has been estimated to be affected by transient salinity (Rengasamy 2006), costing the Australia farming economy approximately \$AUS 1330 million per year (Rengasamy 2002).

Plants have developed numerous mechanisms allowing them to adapt to and tolerate salt stress. Ionic tissue tolerance is one such mechanism. Ion toxicity occurs when normal cellular metabolic processes are disrupted as a consequence of increased ion concentrations.  $Na^+$  ions are the principal cause of ion toxicity caused to cells by salinity stress, resulting in the significant disturbance to cell metabolism, enzymatic activity, and ionic homeostasis (Ahmad *et al.* 2014; Anschütz *et al.* 2014; Shabala *et al.* 2014). To perform well under saline conditions,  $Na^+$  ions should be excluded from shoots, whereas  $K^+$  ions kept at high level ,

thereby maintaining a high cytosolic  $K^+/Na^+$  ratio, especially in leaves (Apse *et al.* 1999; Dubcovsky *et al.* 1996; Gorham *et al.* 1987; Ren *et al.* 2005; Schachtman *et al.* 1989; Serrano *et al.* 2001; Shi *et al.* 2002a). Plants sustain high  $K^+/Na^+$  ratios in the cytosol by expelling  $Na^+$  ions across the plasma membrane and/or by intracellular compartmentalisation of  $Na^+$  ions within the plant vacuole. These processes are crucial for cytosolic  $Na^+$  ion detoxification and for cellular osmotic adjustment, conferring tolerance to salt stress (Blumwald 2000).  $Na^+/H^+$  antiporters exchange protons for  $Na^+$  ions when moving  $Na^+$  across membranes and are particularly active in the vacuoles of plants, algae and fungi. This type of antiporter is also known to exist in animals, yeasts, bacteria and plants (Blumwald 2000). A gene encoding a  $Na^+/H^+$  exchanger (*NHX1*) has been previously identified as encoding a protein important in reducing the accumulation of  $Na^+$  in the cytosol compartmentalizing  $Na^+$  in prevacuolar and vacuolar compartments (Nass *et al.* 1997; Nass *et al.* 1998). The *NHX1* protein is located at the tonoplast and removes  $Na^+$  ions from the cytoplasm. This transporter is energized by the electrochemical potential created by the pumping of  $H^+$  into the vacuole by two proton pumps, vacuolar  $H^+$ -inorganic pyrophosphatase (V-PPase, E.C. 3.6.1.1.) and vacuolar  $H^+$ -ATPase (V-ATPase, E.C. 3.6.1.3) (Gaxiola *et al.* 2007; Sze *et al.* 1999). The *Arabidopsis thaliana*  $Na^+/H^+$  antiporter gene (*AtNHX1*) was the first plant *NHX* homolog to be cloned (Gaxiola *et al.* 1999). There are six isoforms of *AtNHX* in *Arabidopsis* with *AtNHX1* and *AtNHX2* having the greatest transcript distribution between tissues and being highly expressed. *AtNHX3* and *AtNHX4* transcript are exclusively found in flowers and roots (Aharon *et al.* 2003; Yokoi *et al.* 2002). *AtNHX5* and *AtNHX6* are believed to be localized in endosomes and functioning in growth and development, salinity tolerance and vesicle trafficking (Bassil *et al.* 2011a; Qiu 2012). Recently, *NHX1* and *NHX2* were found to transport  $K^+$  into the vacuole giving them a new role in  $K^+$  homeostasis in *Arabidopsis* (Barragan *et al.* 2012; Bassil *et al.* 2011b).

Previous studies have shown improvement in salt tolerance by expressing *NHX1* in a range of salt excluding plants. The pioneering work by Apse et al (1999) has reported the beneficial effect of over-expressing *NHX1* in *Arabidopsis* (Apse *et al.* 1999). Improvement in salinity tolerance was then reported for transgenic plants expressing barley *Hv-NHX2* in *Arabidopsis* (Bayat *et al.* 2011), cotton *GhNHX1* in tobacco plants (Wu *et al.* 2004), *Arabidopsis AtNHX1* in tomato, *Brassica napus* (Zhang *et al.* 2001a; Zhang *et al.* 2001b) and wheat (Xue *et al.* 2004), and *Thellungiella halophila ThNHX1* in *Arabidopsis* (Wu *et al.* 2009). However, all these studies were conducted on salt-sensitive glycophytes whose “default strategy” is Na<sup>+</sup> exclusion from uptake. Will such beneficial effects be observed in “salt includers”, such as barley? In this study we report that merely expressing of *Arabidopsis AtNHX1* in barley has no benefit in increasing plant performance at a whole plant level under saline condition. The possible reasons for the lack of benefit in biomass and other physiological parameters and the implications for breeding salt tolerant plants are discussed.

## **4.2 Materials and Methods**

### **4.2.1 Plant Material and Growth Condition**

The plant material used throughout this study was barley (*Hordeum vulgare* L., cv. Golden Promise). The barley seeds were germinated and grown in 2 L pot filled with University of California (UC) soil mixture (1:1 peat: sand) (Schilling *et al.* 2014). The seeds were watered with reverse osmosis water (RO water) for approximately two weeks until the crop third leaf had fully emerged. Subsequently, NaCl salt was applied for the next five weeks once in two days with 0 mM, 200 mM or 250 mM NaCl. Two *AtNHX1* expressing barley lines and a null segregant were used in the study. For each line there were 6 pots containing 4 plants per pot per treatment. Experiments were conducted between February and April, 2013 in a controlled greenhouse in Adelaide (day/night temperature +25/+10°C, 55-75% relative humidity). Plants were grown under a natural day light.

### 4.2.2 Cloning of *AtNHX1* and generating transgenic lines

The coding sequence of *AtNHX1* (*At5g27150*) was cloned from cDNA by PCR and ligated into g pCR<sup>TM</sup> 8/GW/TOPO<sup>®</sup> TA entry vector. After confirmation of orientation and sequence integrity the transgene was transferred to pMDC32 expression vector using an LR reaction (Gateway Technology, Invitrogen; <http://www.invitrogen.com>) (Jacobs *et al.* 2007). The expression vector was transformed to *Agrobacterium tumefaciens*. Transgenic barley (cv. Golden Promise) expressing *AtNHX1* using the CaMV 35S promoter was successfully generated via *Agrobacterium*-mediated transformation (Jacobs *et al.*, 2007; Singh *et al.*, 1997). The T<sub>3</sub> transgenic barley lines were used for this study. The control plants used for the study were null segregant lines.

### 4.2.3 Genotyping of 35S:*AtNHX1* in barley

Genomic DNA was extracted from leaf tissue of transgenic and null segregants using Edwards, Johnston and Thomson (1991) method. Forward and reverse primers: 5'-ACT CAT AAG CTA CCT ATT ACC G-3' and 5'-TTA CTA AGA TCA GGA GGG TTT CTC-3' were used to amplify a 309 bp amplicon of *AtNHX1* and as a control gene, the *HvVRT2* vernalization gene (GenBank DQ201168) was used and amplified using *HvVRT2*-specific forward primer 5'-CCG AAT GTA CTG CCG TCA TCA CAG-3' and reverse primer 5'-TGG CAG AGG AAA ATA TGC GCT TGA-3', which amplified a fragment of 280 bp in size to determine the presence or absence of the gene of interest. Similar PCR conditions were used for both gene of interest and control gene. ImmoMix<sup>TM</sup> Red (BIOLINE) was used for the PCR (39 cycles of denaturation 94°C for 1 min, annealing 55°C for 1 min and extension 72°C and a final extension of 72°C for 10 min). PCRs contained 2 × reaction of ImmoMix<sup>TM</sup> Red, 10µM of each forward and reverse primers) 1mg/ml BSA, and 0.5 µL of template DNA and ImmoMix Red is based on IMMOLASE<sup>TM</sup> DNA Polymerase. Gel electrophoresis was performed using 2% agarose with 5 µL/100 ml Gel Red to visualise the bands.



#### 4.2.4 Determination of transgene expression

Total RNA was extracted using BIOLINE kit. RNA was diluted to equal concentration of 10 ng/μl and subjected to one step RT-PCR using QIAGEN kit. For the amplification of the *AtNHX1* a forward 5' CCG CAA TTT CGT GAC TAT TAT G 3' and reverse 5' GCT TCG TGG TTT AGG TGA G 3' primer were used with a product size of 309bp. *HvGAP* gene (GenBank EF409629) was used as a positive control (forward primer 5'-GTG AGG CTG GTG CTG ATT ACG-3' and reverse primer 5'-TGG TGC AGC TAG CAT TTG ACA C-3'; product size of 189 bp). The temperature of the one step RT-PCR was programmed at 50°C for 30 min to activate the reverse transcriptase (RT) and 95°C for 15 min to inactivate the RT and denature the cDNA and 35 cycle of denaturation at 94°C for 1 min, annealing 55°C for 1 min and extension 72°C and a final extension of 72°C for 10 min. The RT-PCR reaction mix contained 5× buffer, 10 μM of each forward and reverse primer pairs for the *AtNHX1* and control gene *HvGAP*, water, QIAGEN OneStep RT-PCR Enzyme Mix 2 μL and template RNA (10 ng/μl).

#### 4.2.5 DNA extraction for Southern blot analysis

Plant material was collected in 10 mL tubes, snap frozen in liquid nitrogen and ground to a fine powder by vortexing with three to five sterilised steel balls (Ø 2 mm). The powder was suspended in 1.4 mL of DNA extraction buffer [1 % (v/v) N-Lauroylsarcosine (sigma)-solute to colourless liquid with 50 mg/ml methanol, 100 mM Tris-HCl, 100 mM NaCl, 100 mM EDTA, 2 % (w/v) polyvinyl-pyrrolidone (PVPP, Sigma-Aldrich, Castle Hill, Australia), pH 8.5 with HCl] and thoroughly mixed for 1 min before addition of 1.4 mL of phenol/chloroform/iso-amyl alcohol (25:24:1). After a further 15 min mixing, the samples were centrifuged at 3600 g for 10 min. The aqueous phase was transferred into a 2 mL microcentrifuge tube and DNA precipitated with 100 μL of 3 M sodium acetate and 800 μL of isopropanol. The samples were left for 1 hr at RT before centrifugation at 10,000 g for 10 min.

The supernatant was removed and the pellet washed in 1 mL 70 % (v/v) ethanol. The pellet was air dried and resuspended in 100  $\mu$ L of ddH<sub>2</sub>O containing 10 mg/mL of RNase (Thermo Fisher Scientific Australia Pty Ltd., Scoresby, Australia). The obtained DNA is of high purity and can be used for subsequent Southern Blot analysis.

#### 4.2.6 Southern blot analysis

The DNA samples for the *AtNHX1* lines (*35S:AtNHX1-1* and *35S:AtNHX1-2*), Null segregant and Golden promise (WT) were subjected to Southern blot analysis. The genomic DNA (1  $\mu$ g) were digested with 20 units of *Bam*HI (which cuts once inside the gene *AtNHX1* and in between the gene and the 35S promoter) and *Hind*III (which cuts only once in the vector, at the start of the promoter). An alkaline Southern blot was assembled using a Hybond-N<sup>+</sup> membrane (Amersham) according to manufacturer's instructions. DNA was fixed to the membrane by cross-linking in a UV Stratalinker 2400 (Stratagene). Probe DNA was produced by PCR using primers to amplify fragment of the *hpt* coding sequence. The probe was labelled with <sup>32</sup>P  $\alpha$ -dCTP prior to hybridisation. The blot was imaged using a Typhoon 9200 Phosphorimager (Amersham).

#### 4.2.7 Biomass Measurements

The fresh weight of the above ground biomass was recorded at the end of the study period. The dry weight of each sample plants was measured, after drying the plants at 65<sup>0</sup>C for 72 hrs in an oven. As an indicator of biomass, plant height was recorded at the end of the study period.

#### 4.2.8 Na<sup>+</sup> and K<sup>+</sup> content of the leaf

At the end of the salt application, the third leaf of barley plants was removed and oven dried as described above. A leaf aliquot of 0.3 g was then digested in 1% nitric acid by heating at 85 <sup>0</sup>C for 4 hrs. on a hot block (Environmental Express, Mt Pleasant, SC, USA). The solution in the digested leaf was diluted with milliQ water in a dilution factor of 1:20. The Na<sup>+</sup> and K<sup>+</sup>

content were measured by a flame photometer model 420 (Sherwood Scientific LTD, Cambridge, UK).

#### 4.2.9 Gas exchange and water relations

The stomatal conductance ( $G_s$ ) was measured from the third leaf after the culmination of the NaCl treatment on a sunny day during the day times when the plants are fully transpiring. For measuring the  $G_s$ , leaf porometer Model SC-1 (Deacagon device INC, WA, Australia) was used. The measurements were taken consistently from one-third from the leaf tip of the third leaf in all samples. The relative water content (RWC) was calculated from the fresh and dry weight of the plants by using the formula:

$$RWC = \frac{FW - DW \times 100}{FW}$$


#### 4.2.10 Statistical analysis

The data was analysed using student's t-test method and the significant level of mean difference was compared at probability value of  $P \leq 0.05$  (\*),  $P \leq 0.01$  (\*\*) and  $P \leq 0.001$  (\*\*\*) using a statistical software SPSS version 22 (IBM support portal, USA).


## 4.3 Results

### 4.3.1 Growth response of barley plants expressing *AtNHX1*


The presence of the *AtNHX1* transgene in the transgenic barley was confirmed by PCR. PCR analysis showed that the *AtNHX1* gene was present in *35S:AtNHX1-1* and *35S:AtNHX1-2* lines and absent in both the null and Golden Promise (GP) (Fig. 4.1a). RT-PCR was performed to confirm that the transgene was expressed in the transgenic lines and not in either of the nulls or the wild type (Fig. 4.1b). Southern blot analysis was done to check the number of inserts of *AtNHX1* in the transgenic lines. *35S:AtNHX1-1* had a single insert while *35S:AtNHX1-2* had three inserts. Both the null line and Golden Promise were found to have no insert (Fig. 4.2). The two barley *AtNHX1* expressing lines, a null segregant and Golden Promise were treated with 200 mM and 250 mM NaCl for five weeks after seedling establishment. *AtNHX1* expressing lines did not show any obvious phenotype and did not gain any advantage in growth and development compared to the wild type or null barley plants under both control and saline conditions (Fig. 4.3). The biomass of all plants was significantly reduced after the addition of salt. Moreover, the fresh weight of *AtNHX1-2* line was 15% lower compared to the null segregant (significant at  $P \leq 0.01$ ). The other transgenic line didn't show significant (at  $P \leq 0.05$ ) reduction in fresh weight compared to the null segregants (Fig. 4.4a). No significant change in dry weight was observed in transgenic lines (Fig. 4.4b). Taken together, these results suggest that expressing *AtNHX1* in barley does not improve plant phenotype under saline conditions.




**Figure 4.1** Molecular analysis of *35S:AtNHX1* lines in barley (a) Confirmation of transgene presence by PCR. GP (Golden Promise), L1 (*35S:AtNHX1-1*), L2 (*35S:AtNHX1-2*), N (null line), (-) negative control. (b) Confirmation of transgene expression in transgenic lines by RT-PCR. A multiplex reaction using primers for *AtNHX1* (upper band) and the control gene *Hv-GAP* (lower band) was performed.



**Figure 4.2** Southern blot analysis of two transgenic lines, a null line and Golden Promise. The Genomic DNA was digested with BamHI and HindIII restriction enzymes and probed by hygromycin DNA. The lanes 1-2 and 10-11 correspond to Golden Promise. Lanes 3-4 and 12-13 correspond to null line. Lanes 5-6 and 14-15 correspond to *35S:AtNHX1-2*. Lanes 7-8 and 16-17 correspond to *35S:AtNHX1-1* with BamHI and HindIII digestion, respectively. Lane 9 and 18 are positive controls.



**Figure 4.3** Growth and development of three barley *35S:AtNHX1* lines and non-transgenic control cultivar Golden promise (GP) treated with 0 mM, 200 mM NaCl and 250 mM NaCl for five weeks after seedling establishment. *35S:AtNHX1-1* corresponds to line L1, *35S:AtNHX1-2* to L2, and GP to Golden Promise.



**Figure 4.4** Fresh weight (a) and dry weight (b) of two barley *35S:AtNHX1* lines and the null line. Results are mean  $\pm$  SE (n=21-24). Statistical significance designated as  $P \leq 0.05$  (\*),  $P \leq 0.01$  (\*\*) and  $P \leq 0.001$  (\*\*\*)

### **4.3.2 Expressing *AtNHX1* in barley gives no advantage to either enhanced gas exchange or relative water content**

The stomatal conductance of all lines tested was reduced by approximately 60 to 70% after treatment with NaCl. Expressing *AtNHX1* in barley did not result in improvements in stomatal conductance under salt stress (Fig. 4.5a). Relative water content of all salt grown lines was significantly reduced when compared to those grown in the absence of salt. The *35S:AtNHX1-2* line showed a 4% reduction in RWC (significant at  $P \leq 0.001$ ) compared to the null segregant. The *35S:AtNHX1-1* line has non-significant difference with the null segregant. Taken together, these results suggest the lack of beneficial, or even some detrimental (as in line *35S:AtNHX1-2*) effects of *AtNHX1* expression for gas and water relations in barley (Fig. 4.5a & b).


### **4.3.3. Expressing *AtNHX1* in barley has not resulted in increased chlorophyll content**

There was no significant difference in chlorophyll content in *35S:AtNHX1-1* when compared to the null line. In *35S:AtNHX1-2* line, a small but significant (at  $P \leq 0.05$ ) decline in chlorophyll content was observed between as compared with the null (Fig. 4.6). This suggests that chlorophyll content could not be positively influenced by expressing *AtNHX1* in barley.


### **4.3.4 Plants expressing *AtNHX1* showed increased shoot $\text{Na}^+$ and $\text{K}^+$ accumulation**

Leaf  $\text{Na}^+$  accumulation increase substantially 3.5-4.5 fold higher in all lines after the addition of NaCl. The *35S:AtNHX1-2* line showed 22% increase in leaf  $\text{Na}^+$  accumulation compared to the null segregant (significant at  $P \leq 0.001$ ). The other transgenic line *35S:AtNHX1-1* also showed slight (14%) but not significant increase in leaf  $\text{Na}^+$  content (Fig. 4.7a). Similarly,  $\text{K}^+$  accumulation in the shoot was approximately 30% higher (significant at  $P \leq 0.001$ ) for the *35S:AtNHX1-2* line compared to the null segregant, and marginally higher (8%; not significant at  $P \leq 0.05$ ) shoot  $\text{K}^+$  accumulation was measured in *35S:AtNHX1-1* (Fig. 4.7b).






**Figure 4.5** (a) Stomatal conductance of two barley *35S:AtNHX1* lines and null lines. Results are the mean  $\pm$  SE (n=21-24) (b) Relative water content of two barley *35S:AtNHX1* lines and a null. Results are mean  $\pm$  SE (n=21-24). Statistical significance designated as  $P \leq 0.05$  (\*),  $P \leq 0.01$  (\*\*) and  $P \leq 0.001$  (\*\*\*)



**Figure 4.6** Chlorophyll content (arb. unit) of two *35S:AtNHX1* barley lines and a null line were exposed to 0 mM, 200mM and 250mM NaCl for five weeks after seedling establishment. Results are the mean  $\pm$  SE (n=21-24). Statistical significance designated as  $P \leq 0.05$  (\*),  $P \leq 0.01$  (\*\*) and  $P \leq 0.001$  (\*\*\*)



**Figure 4.7** Leaf Na<sup>+</sup> (a) and K<sup>+</sup> (b) content of two 35S:AtNHX1 barley lines and null (used as control) exposed to 200mM and 250mM NaCl for five weeks after seedling establishment. Mean  $\pm$  SE (n=21-24). Statistical significance designated as  $P \leq 0.05$  (\*),  $P \leq 0.01$  (\*\*) and  $P \leq 0.001$  (\*\*\*)

## 4.4 Discussion

### 4.4.1 Expressing AtNHX1 alone in barley has no advantage for biomass

The biomass of the *AtNHX1* expressing barley lines has shown non-significant change compare to the null segregants, suggesting that the expression of *AtNHX1* in has no beneficial effect in increasing biomass. Moreover, *AtNHX1-2* line even showed a small but significant reduction in fresh weight under 250 mM salt treatment (Fig. 4.4a & b). Several possible reasons may explain this lack of beneficial effects. One of them is a possible misfolding of the *AtNHX1* protein which would result in an inactive protein and no improvement in phenotype. Another reason might be the mistargeting of the protein to other membranes, where it is not usually associated with. Constitutive over-expression of genes encoding membrane proteins is known to often result in a large amount of protein produced, leading that protein appearing on other membranes. For example over-expression of *AVP1* can result in the vacuolar H<sup>+</sup>-PPase

being targeted to the plasma membrane (Gaxiola *et al.* 2012; Li *et al.* 2005). Thirdly, the lack of phenotype may be due to NHX1 protein not undergoing the required post-translational modifications, such as by a kinase, to activate its transporter capabilities (Neuhaus *et al.* 2014; Weinl *et al.* 2009). It has been reported that NHX proteins may be targeted by the CBL-CIPK calcium signalling pathway, similar to the regulation of AtSOS1 by the kinase AtSOS2 (AtCIPK24) (Qiu *et al.* 2004).

Apart from issues with the production of an Arabidopsis protein in barley there are other physiological reasons as to why no phenotypic difference was observed between the transgenic lines and nulls. First, the ‘pump and leak concept’ explained for plasma membrane of algal cell (Brummer *et al.* 1985) also works for plasmalemma and tonoplast of higher plant (Hedrich *et al.* 2011). Vacuolar Na<sup>+</sup> may back-leak to the cytosol via Na<sup>+</sup> permeable SV (slow activating) and FV (Fast activating) channels, and it was shown that reduced activity of these channels confer salt tolerance in halophyte species (Bonales-Alatorre *et al.* 2013b; Hedrich *et al.* 1987). SV channels conduct both mono- and divalent cations indiscriminately based on Ca<sup>2+</sup> and voltage dependence and FV channels conduct monovalent cations including K<sup>+</sup> (Pottosin *et al.* 2014). It was suggested earlier that, to achieve efficient vacuolar Na<sup>+</sup> sequestration, increased NHX activity should be accompanied by efficient control of SV and FV channels permeability (Bonales-Alatorre *et al.* 2013b; Shabala 2013). This may be not the case in this study. Second, the proton pumps V-ATPase and V-PPiase are responsible for developing proton motive force to move ions into the vacuole (Gaxiola *et al.* 2007; Sze *et al.* 1999), and their high activity is essential for efficient vascular sequestration of Na<sup>+</sup> by NHX (Shabala 2013). Thus, it may appear that in barley the intrinsic tonoplast H<sup>+</sup>-pumping activity is not sufficient to drive any “extra” NHX, and concurrent expression of NHX transporters and H<sup>+</sup> pumps is required to confer salinity tolerance. Third, vacuolar H<sup>+</sup>-ATPase pumping require ATP and, even plants may have enough functional AHA proteins encoding this

pumping, they may be simply short of ATP to make them functional. Forth, the reduced  $K^+$  pool in the cytosol due to its sequestration in the vacuole would seek compensation by production of osmolytes to prevent dehydration. However, the mitigating effect of osmolytes for loss of  $K^+$  from the cytosol in tolerant barley genotypes is less pronounced (Chen *et al.* 2007a) and the genetic background used for this study is a comparatively tolerant barley cultivar (Adem *et al.* 2014).

Taken together, it appears that transgenic barley expressing *AtNHX1* requires pyramiding with genes that will energize the tonoplast membrane. It was shown earlier that the co-expression of *Suaeda salsa*  $Na^+/H^+$  antiporter (*SsNHX1*) and *AVP1* has shown higher salinity tolerance in rice than the expression of *SsNHX1* alone (Zhao *et al.* 2006) and required a kinase gene to assist in the activity of the NHX protein. Also, its deemed necessary considering to control back leak of  $Na^+$  in the positive membrane potential sensing  $Na^+$  permeable SV and FV channels (Shabala 2013).

#### **4.4.2 Barley plants expressing *AtNHX1* alone has shown no improvement in water relation and gas exchange characteristics**

Stomatal conductance is one of physiological trait that is considered as a yield determinant (Lu *et al.* 1998; Richards 2000). Control of stomatal conductance to balance for optimum transpiration rate and water retention is of paramount importance for a plant growth under salt stress (Munns 2002). Cognizant to these facts, there was no increase in stomatal conductance in *35S:AtNHX1-1* line and even there was a significant reduction  $P \leq 0.001$  in *35S:AtNHX1-2* line compared to the null segregant. Similar reduction was observed in the fresh weight and RWC. In previous experiment Golden Promise showed a better water retention under salt stress (Adem *et al.* 2014), however, in this experiment it was not possible to improve water retention further for Golden Promise. Presumably a leaky system could be assumed for the  $Na^+$  and  $K^+$  ions which are used as osmoticum for maintain cell turgour, stomatal aperture and

gas exchange and ultimately water relation of the plant where the factors are crucial for adaptation under water stress (Anschütz *et al.* 2014; Mengel *et al.* 1982). Therefore, for obtaining improvement in water relation and gas exchange using *AtNHX1* expression in barley has to take into consideration all factors involved in efficient sequestration of ion and proper feedback mechanisms in ion homeostasis which are discussed in this paper.

#### **4.4.3 Sodium and Potassium accumulated in the leaf of barley plants expressing *AtNHX1***

The Na<sup>+</sup> content in the leaf of the barley plants expressing *AtNHX1* has increased significantly ( $P \leq 0.001$ ) as shown in *35S:AtNHX1-2* line in comparison to null segregants. The two possible scenarios may be higher pumping of Na<sup>+</sup> into the vacuole and hence, higher Na<sup>+</sup> in the shoot or the pleiotropic alterations of native gene expression in barley due to the expression of the Arabidopsis transgene which may have unforeseen effects on abundance and activity of proteins involved in Na<sup>+</sup> transport in a plant. Alterations in native gene expression related to ion transport were observed in transgenic rice, after cell type specific expression of *AtHKT1;1* in root cortical cells, which were hypothesized to alter Na<sup>+</sup> transport through the transgenic plants (Plett *et al.* 2010a). Mis-expression of *AtNHX1* in barley may have altered genes encoding proteins which result in a higher translocation of Na<sup>+</sup> to the shoot. Similarly, leaf K<sup>+</sup> content of the *35S:AtNHX1-2* was shown to be higher compared to the null segregants. This may be attributed to the K<sup>+</sup>/H<sup>+</sup> transport activity of the *NHX* protein reported recently (Barragan *et al.* 2012; Bassil *et al.* 2011b; Leidi *et al.* 2010; Rodriguez-Rosales *et al.* 2008). The lower K<sup>+</sup> level in the cytosol due to its transport to the vacuole may be one reason for lack of phenotype in the barley plants expressing *AtNHX1*. The *AtNHX1* is not only a Na<sup>+</sup> transporter but also K<sup>+</sup> as cited above. The *AtNHX1* transports K<sup>+</sup> to the vacuole reducing the cytosol K<sup>+</sup> pool which is detrimental as a result counteracting the beneficial effect of the protein, therefore, no benefit in performance under salt stress. In addition, the salt induced

repression of *HAK* as shown in *LeHAK5* repression of expression under salt growth condition (Nieves-Cordones *et al.* 2008) which may be considered as other factor for salt tolerance by limiting  $K^+$  uptake under salinity stress.

Taken together, the lack of phenotype might be hypothesized due to a number of reasons: (i) low level of activity of vacuolar  $H^+$ -inorganic pyrophosphatase (V-PPase, E.C. 3.6.1.1.) and vacuolar  $H^+$ -ATPase (V-ATPase, E.C. 3.6.1.3) resulting in an insufficient proton gradient required for vacuolar  $Na^+/H^+$  exchanger; (ii) the inability of transgenic plants to prevent a passive leak of sodium via  $Na^+$  permeable slow activating (SV) and fast activating (FV) channels; (iii) insufficient ATP pool to support  $H^+$  pumping activity; and (iv) the fact that the *NHX1* protein might not been properly folded, inactive or incorrectly targeted. Furthermore, the above mentioned notions can be strengthened by observing the contribution of ectopically expressed *AtNHX1* to endogenous  $Na^+/H^+$  and  $K^+/H^+$  antiport activities in the vacuole of control plants, which could be intrinsically high considering that barley is a salt inculder and cation/proton exchange activity in the transgenic in comparison to control plants.

## **Chapter 5      Expressing *Arabidopsis thaliana* V-ATPase subunit C in barley (*Hordeum vulgare* L.) improves plant performance under saline condition by enabling better osmotic adjustment <sup>4</sup>**

Getnet D. Adem<sup>1</sup>, Stuart J. Roy<sup>2,3</sup>, Meixue Zhou<sup>1</sup>, John Bowman<sup>1</sup> and Sergey Shabala<sup>1\*</sup>

<sup>1</sup>School of Land and Food, University of Tasmania, Private Bag 54, Hobart, Tas 7001, Australia; <sup>2</sup>Australian Centre for Plant Functional Genomics, Private Mail Bag 1, Glen Osmond, SA 5064, Australia; <sup>3</sup>School of Agriculture, Food and Wine, University of Adelaide, Private Mail Bag 1, Glen Osmond, SA 5064, Australia

\*Corresponding author; email: Sergey.Shabala@utas.edu.au

### **Summary**

Salinity is a global problem affecting agriculture resulting in an estimated \$27Bln per annum loss revenue. Overexpressing of vacuolar ATPase subunits has been shown to be beneficial in improving plant performance under saline conditions. Most studies, however, have not shown if there is an improvement on grain yield, and/or investigated the physiological mechanisms behind the improvement in plant growth. In this study, we constitutively expressed *Arabidopsis Vacuolar ATPase subunit C* (*AtVHA-C*) in barley. Transgenic plants were assessed for agronomical and physiological characteristics, such as dry biomass, leaf pigment content, stomatal conductance, grain yield, and leaf Na<sup>+</sup> and K<sup>+</sup> content, when grown with either 0 or 300 mM NaCl. When compared to non-transformed barley, the *AtVHA-C* expressing barley lines have a smaller reduction in both biomass and grain yield. The ability of the transgenic lines to maintain higher stomata conductance, resulting from an increase in

---

<sup>4</sup> This paper has been submitted and is currently under review in *Plant Biotechnology Journal*

$\text{Na}^+$  and  $\text{K}^+$  accumulation in the leaf which could be used for osmotic adjustment rather than the synthesis of energy expensive organic osmolytes.

**Key words:** vacuolar sequestration; osmotic adjustment; salinity stress tolerance; sodium; potassium; organic osmolytes

## 5.1 Introduction

With the world population estimated to exceed 9.3 billion by 2050, there is a shift towards agricultural production systems using of marginal saline lands (Panta *et al.* 2014; Shabala 2013). The problem is further aggravated by the reduction in precipitation in subtropical areas, which results in farmers irrigation with brackish and low quality water, as the good quality water is reserved for drinking (Barrett-Lennard *et al.* 2010; Munns *et al.* 2008). The estimated cost of salinity-induced loss to crop production is believed to be in excess of \$27 Bln per year (Qadir *et al.* 2014). Improving performance of crop plants so that they produce a better yield under high saline conditions is of paramount importance for using these marginal lands (Munns 2002; Roy *et al.* 2013).

Soil salinity reduces the amount of available water requires plants to make osmotic adjustments to grow in these conditions. Inorganic ions such as  $\text{K}^+$ ,  $\text{Na}^+$  and  $\text{Cl}^-$  play an essential role in this process, contributing to 80-95% of the cell sap osmotic potential in halophytes (Glenn *et al.* 1999) and to 50-70% in glycophytes, such as barley and wheat (Chen *et al.* 2007b; Cuin *et al.* 2010). However, excess  $\text{Na}^+$  in the cytosol is toxic to cellular processes, regardless of whether plants are halophytes or glycophytes (Flowers *et al.* 2008). Therefore for  $\text{Na}^+$  to be an effective osmolyte it must be sequestered in the vacuole. Three components are essential for the vacuolar storage of  $\text{Na}^+$ : establishment of the electrochemical gradient necessary to pump  $\text{Na}^+$  into the vacuole;  $\text{Na}^+$  transport across the tonoplast into the vacuole; and  $\text{Na}^+$  retention within the vacuole (Shabala 2013). Tonoplast  $\text{Na}^+/\text{H}^+$  antiporters such as members of the  $\text{Na}^+/\text{H}^+$  exchanger (NHX) family have been shown to be important in



the loading of  $\text{Na}^+$  into the vacuole (Apse *et al.* 2007; Barkla *et al.* 1995; Blumwald *et al.* 1985; Gaxiola *et al.* 1999). There are numerous NHX proteins which are expressed in different tissue- and stress-dependent manner (Rodriguez-Rosales *et al.* 2008). Proteins such as NHX rely on a proton gradient between the vacuole and the cytoplasm which is generated by two types of tonoplast localized proton pumps, namely vacuolar ATPase (VHA) (Krebs *et al.* 2010; Vera-Estrella *et al.* 1999; Wang *et al.* 2001) and vacuolar pyrophosphatase (V-PPase) (Guo *et al.* 2006; Parks *et al.* 2002; Vera-Estrella *et al.* 2005). Lastly, to retain sequestered  $\text{Na}^+$  in the vacuole, preventing it from leaking back into the cytosol efficient control of expression and activity of sodium permeable slow (SV) and fast (FV) vacuolar channels is required (Bonales-Alatorre *et al.* 2013a; Bonales-Alatorre *et al.* 2013b; Pantoja *et al.* 1989; Shabala *et al.* 2011a).

Many studies investigating whether improvements in vacuolar  $\text{Na}^+$  sequestration can improve salinity tolerance, usually overexpression a single NHX gene. Results ranged from highly positive (Apse *et al.* 1999; Bayat *et al.* 2011; Zhang *et al.* 2001a; Zhang *et al.* 2001b) to no yield improvement or even detrimental effects (Adem *et al.* 2015). A better approach may be in manipulating the expression of genes encoding proteins which set up the necessary vacuolar conditions for effective  $\text{Na}^+$  sequestration, such as manipulation of the expression of V-ATPases and V-PPases.

Overexpression of the Arabidopsis vacuolar  $\text{H}^+$ -PPase (AVPI) has conferred both salinity and drought tolerance in Arabidopsis (Gaxiola *et al.* 2002; Gaxiola *et al.* 2001) and overexpression of  $\text{H}^+$ -PPase from *Suaeda salsa* improved salinity tolerance in Arabidopsis (Guo *et al.* 2006). Consistent with these observations, Zhao *et al.*, (2006) showed that the co-expression of the *Suaeda salsa*  $\text{Na}^+/\text{H}^+$  antiporter (*SsNHX1*) and AVPI conferred higher salinity tolerance in rice than the expression of *SsNHX1* alone.

H<sup>+</sup>-PPases are often considered a supporting mechanism to V-ATPases in the acidification of the vacuole (Maeshima 2000). Hence, investigating whether greater improvements in vacuolar storage of Na<sup>+</sup> and in salinity tolerance can be made by manipulating the abundance of V-ATPases is important.

Unlike V-PPases, vacuolar H<sup>+</sup>-V-ATPase (*VHA*) are made of several subunits, encoded by different genes. This makes overexpressing the entire protein difficult. Studies have investigated, however, whether manipulation of the expression of genes encoding specific V-ATPase subunits can improve plant growth under saline conditions. Expressing the V-ATPase c1 subunit from the halophyte grass *Spartina alterniflora* improved rice growth under saline conditions (Baisakh *et al.* 2012). However, the beneficial effect was reported only for vegetative plant growth but not grain yield. Interestingly, the observed effect was explained by the early closure of the leaf stoma and reduced stomata density (Baisakh *et al.* 2012) during salt stress, which is not consistent with the transgenic plants increasing their biomass when they have reduced capacity to assimilate CO<sub>2</sub>. More recently, genes encoding individual V-ATPase subunits in wheat line were expressed in *Arabidopsis* (He *et al.* 2014). While improvements in the salt tolerance of seedling transgenic *Arabidopsis* plants were observed, no insights into the mechanisms behind the improved tolerance were reported (He *et al.* 2014). Similar results were reported when wheat *V-ATPase subunit B* (Wang *et al.* 2011) and *subunit E* (Zhao *et al.* 2009) were expressed in *Arabidopsis* and when the *V-ATPase subunit c* (*VHA-c*) from *Limonium bicolor* was expressed in tobacco (Xu *et al.* 2011).

All the above examples come from salt-sensitive species (rice; *Arabidopsis*; tobacco) that predominately rely on sodium exclusion to achieve their tolerance (Munns *et al.* 2008). There are other species, such as barley, which will still predominately exclude the majority of the Na<sup>+</sup> in the soil from the shoot and are better able at accumulating higher concentrations of Na<sup>+</sup> in the shoot, relying on Na<sup>+</sup> to achieve their osmotic adjustment under saline conditions

(Shabala *et al.* 2010). The questions remains as to what extent overexpression of *V-ATPase* subunits may enhance salinity stress tolerance in barley, and the mechanisms behind any beneficial effect observed.

In this study, we generated lines of barley expressing the *Arabidopsis V-ATPase subunit C* (*VHA-C*) and assessed their physiological and agronomical performance under saline conditions. *V-ATPase* subunit C was chosen as this is on peripheral of the V1 complex, which has a function in ATP hydrolysis (Perez-Sayans *et al.* 2012; Sze *et al.* 2002), and can directly enhance the activity of the *V-ATPase*. Barley lines expressing the *AtVHA-C* gene were found to have smaller salinity driven reductions both biomass and grain yield compared to non-transformed wild type plants. These beneficial effects may be linked to the improved ability of the transgenic lines to maintain higher stomata conductance, resulting from increased  $\text{Na}^+$  and  $\text{K}^+$  accumulation in the leaf which contributed to osmotic adjustment and made plant less reliant on *de novo* synthesis of organic osmolytes.

## **5.2 Experimental procedures**

### **5.2.1 Generating transgenic barley plants expressing *VHA-C* (*Arabidopsis V-ATPase Subunit C*)**

The coding sequence of *V-ATPase subunit C* (*At1g12840*) was amplified from the cDNA of *Arabidopsis thaliana* (ecotype Col-0) using primer pairs, 5' -AGA GAC TCG TAA ACA AGA G -3' and 5' -CAG CCA TGG CTC CTG CA -3' and a high fidelity DNA polymerase. The amplified sequence was introduced into the *pCR8/GW/TOPO TA* Gateway enabled entry vector (Invitrogen). The *V-ATPase C* was then recombined into a *pMDC32* destination vector Gateway LR<sup>®</sup> recombination reaction (Invitrogen, Carlsbad, CA, USA) (Curtis *et al.* 2003), 3' of a *CaMV 35S* promoter. The *VHA-C pMDC32* vector was transformed into the barley, *Hordium vulgare*, cultivar WI4330 (kindly supplied by the University of Adelaide Barley Breeders. the by the *Agrobacterium tumefaciens*- mediated

transformation method using the ACPFG (University of Adelaide) transformation facility. After antibiotic selection, the transformed plantlets were regenerated on soil (Jacobs *et al.* 2007; Singh *et al.* 1997) to produce T<sub>1</sub> seed. After one round of seed multiplication the T<sub>2</sub> plants were grown for experiment in the University of Tasmania.

### 5.2.2 Plant growth conditions

Four *VHA-C* expressing (35S:*VHA-C*) T<sub>2</sub> lines, described here as OE1, OE2, OE3 and OE4, and one non-transfected WI4330 were grown on 2L pot containing potting mix. The potting mix was composed of 70% composted pine bark; 20% coarse sand; 10% sphagnum peat; Limil at 1.8 kg/m<sup>3</sup>, dolomite at 1.8 kg/m<sup>3</sup>). The plant nutrient balance was maintained by adding the slow release Osmocote Plus™ fertilizer (at 6 kg/m<sup>3</sup>), plus ferrous sulphate (at 500 g/m<sup>3</sup>) (Bonales-Alatorre *et al.* 2013a). The plants were grown in a glasshouse with regulated temperature (24°C-20°C day/15°C night) for approximately 2 weeks (until the 3<sup>rd</sup> leaf stage) until the seedling established. Then salinity treatment was administered by adding 300mM NaCl with irrigation water until maturity. The experiment was conducted between January and April 2014. Plants were grown in 4L pots, with three plants per pot and 4 to 5 pots per treatment/genotype.

### 5.2.3 DNA extraction and PCR analysis

Barley genomic DNA from the above lines was extracted using (Edwards *et al.* 1991) method. First, the presence of DNA and the PCR condition was confirmed using a control gene, *Hv-VRT2* vernalization gene (GenBank DQ201168) which was amplified using *Hv-VRT2*-specific forward primer 5'-CCG AAT GTA CTG CCG TCA TCA CAG-3' and reverse primer 5'-TGG CAG AGG AAA ATA TGC GCT TGA-3', which amplified a fragment of 280 bp in size. The *VHA-C* gene was detected using a- forward primer 5'-AGA GAC TCG TAA ACA AGA G -3' and reverse primer 5'-CAG CCA TGG CTC CTG CA -3', which amplified a fragment of 299 bp in size. PCRs contained 2 × reaction of ImmoMix™ (BIOLINE), forward

and reverse primers (10  $\mu$ M each), BSA (1 mg/ml), MilliQ water and 0.5  $\mu$ L of template DNA. The PCR conditions used were similar for both the *VRT2* and the *VHA-C* (95°C for 10 min, followed by 1 min at 94°C, 1 min at 55°C and 1min at 72°C, repeated for 35 cycles, with a final extension of 72°C for 10min). PCR products were visualised by gel electrophoresis using a 2% agarose with 5 $\mu$ L/100ml Gel Red Nucleic acid stain (BIOTIUM) and the image was taken using gel doc (Molecular imager, Gel Doc<sup>TM</sup> imaging system, BIORAD).

#### 5.2.4 RNA extraction and gene expression analysis

Total RNA was extracted from leaf tissue using BIOLINE plant Isolate II RNA kit ([www.bioline.com/isolate](http://www.bioline.com/isolate)). The RNA was diluted to 10 ng/ $\mu$ l and subjected to a RT-PCR with a master mix containing components of 5  $\times$  buffer, dNTP (10 mM), water and enzyme mix (*Taq* polymerase and reverse transcriptase), forward and reverse primers for *VHA-C* and *HvGAP* (*VHA-C* gene: forward primer 5'-AGA GAC TCG TAA ACA AGA G -3' and reverse primer 5'-CAG CCA TGG CTC CTG CA -3'; and *Hv-GAP*: forward primer 5'-GTG AGG CTG GTG CTG ATT ACG-3' and reverse primer 5'-TGG TGC AGC TAG CAT TTG ACA C-3') and a template RNA of 10 ng/ $\mu$ l. Reverse transcription was performed at 50°C for 30 min, followed by PCR (initial PCR activation 95°C for 15 min, then 35 cycles of 94°C for 1min, of 55°C for 1 min and 72°C for 1min, before a final extension 72°C for 10min). The PCR product was visualized by gel electrophoresis using a 2% agarose gel containing 5  $\mu$ L/100ml of Gel Red Nucleic acid stain (BIOTIUM) using gel doc (Molecular imager, Gel Doc<sup>TM</sup> imaging system, BIORAD).

#### 5.2.5 Biomass and grain yield

As grain yield was required, only the dry weight of the plants (not fresh weights) was recorded at the end of experiment, alongside with plant height. The number of seeds per plant and their total weight were recorded. The sample size for biomass and yield data for the salt treated plants range from 8 to 15. The number of chlorotic and necrotic leaves per plant was

also recorded after 5 weeks of salt treatment. For chlorotic leaves, the partially yellow leaves per plant was counted and presented as chlorotic leaves. For necrotic leaves, the dead leaves per plant was counted and presented as necrotic leaves.

### **5.2.6 Pigment content and stomatal conductance**

Relative leaf chlorophyll content was measured using Minolta Chlorophyll Meter SPAD-502 (Konica Minolta, Osaka, Japan). Measurements were taken from the third true leaf, at a position about one quarter of the length of the leaf from the leaf tip. The same leaf/position was used to measure stomatal conductance (Gs) from control and salt treated plants. These measurements were conducted using Decagon leaf porometer (Decagon Devices Inc., WA, Australia), under constant light conditions (artificial light of  $150 \mu\text{mol m}^{-2} \text{s}^{-1}$ ) between 11 am and 3 pm.

### **5.2.7 Na<sup>+</sup> and K<sup>+</sup> content**

The third leaf from both salt treated and control plants were excised after five weeks of salt treatment for Na<sup>+</sup> and K<sup>+</sup> ion content measurement. Leaves were dried in the oven at 65°C for 72 hrs. A subsample of a known weight (about 0.2 g) was then digested in 5ml of 65-70 % of HNO<sub>3</sub> and 2ml of 30% of H<sub>2</sub>O<sub>2</sub> using a microwave digester (MDS-2000, CEM corporation, Matthews, NC, USA). The digested samples were diluted 1:5 using distilled water. The Na<sup>+</sup> and K<sup>+</sup> ion content was measured using a flame photometer (MODEL PFP7-Flame photometer, JENWAY, Bibby Scientific Ltd, UK). The concentration of the ion content was calculated as mmol kg<sup>-1</sup> DW taking aliquot weight and dilution factors into consideration.

### **5.2.8 Sap osmolality measurements and calculation of contribution of inorganic ions towards leaf osmotic adjustment**

After five weeks of treatment with 350 mM NaCl, third leaf of the barley transgenic and WT plants harvested and stored at -20°C. The leaf sap was extracted using freeze-thaw method (Tomos *et al.* 1984) and osmolality was measured using the vapour pressure osmometer

(Vapro, Wescor Inc. Logan, Utah, USA). The sample size for both control and salt treated plants was  $n=6$ . Measured  $\text{Na}^+$  and  $\text{K}^+$  concentrations were then used to calculate their relative contribution towards overall sap osmotic potential. Contribution of  $\text{Cl}^-$  was estimated as 1.2 of that for  $\text{Na}^+$ , as previously described in the literature (James et al., 2006; Puniran-Hartley et al., 2014).


### 5.2.9 Statistical analysis

The data was analysed using student's t-test method and the significant level of mean difference was compared at probability value of 0.05 (\*), 0.01 (\*\*) and 0.001 (\*\*\*) using a statistical software SPSS version 22 (IBM support portal, USA).

## 5.3 Results

### 5.3.1 Generation of barley plants expressing *V-ATPase Subunit C*

Transgenic barley (cv. WI4330, kindly supplied by the University of Adelaide Barley Breeding Program) expressing the *V-ATPase subunit C* (*VHA-C*) was generated using an agrobacterium mediated transformation (Jacobs *et al.* 2007; Tingay *et al.* 1997). Four *AtVHA-C* expressing lines, designated as OE1-OE4 (*35S:AtVHA-C-1* to *35S:AtVHA-C-4*), and one non-transformed WI4330 line (WT) were included in this study. The OE4 line did not show consistent phenotype throughout all experiments, most likely due to a detrimental effect from the transformation process, and was therefore omitted from analysis. The presence of the *VHA-C* gene in the genome of transgenic lines and its absence in the WT were confirmed using PCR on genomic DNA extracted from the leaf (Fig. 5.1a). The expression of the *VHA-C* gene in transgenic lines and the absence of its expression in the WT were confirmed by RT-PCR (Fig. 5.1b).




**Figure 5.1** Genotyping *AtVHA-C* transgenic barley. (a) Detection of transgene presence by PCR using specific primers for *AtVHA-C* and *Hv-VRT2* (internal control); (b) Expression analysis of the *AtVHA-C* gene using RT-PCR with specific primers for the gene and *HvGAP* specific primers (internal control). Lanes 35S:*AtVHA-C*-1, 2, 3, & 4 are transgenic *AtVHA-C* barley lines, WT (non-transformed barley cv. WI4330), (-) negative control (water).



### **5.3.2 Transgenic barley expressing *V-ATPase subunit C* has improved overall salinity stress tolerance and grain yield**

The *AtVHA-C* expressing lines (OE) had larger seed size and were significantly ( $P \leq 0.001$ ) heavier compared to WT barley plants after growing under 300 mM NaCl stress (Fig. 5.2; Table 5.2). The better performance of the OE lines was also reflected in plant phenotype (Fig. 5.3), with the transgenic lines producing more tillers and remaining green, while the WT plants showed signs of severe senescence. While no significant (at  $P \leq 0.05$ ) difference was found between the dry weight (DW) of OE lines and WT under control conditions (0 mM NaCl; Fig. 5.4a), and an slight increase in the DW of transgenic plants under salt stressed conditions (300 mM NaCl; Fig 5.4b), the relative DW reduction under saline conditions was significantly lower in all transgenic lines compared with WT (Fig. 5.4c). The relative DW of salt-treated overexpressing lines ranged within 43 to 47% of control vs only 35% in WT (Fig. 5.4c). While there was little difference in plant height in the 0 mM treatment (Fig 5.4d), all saline-grown OE lines were much taller compared with WT (Fig. 4e; significant at  $P \leq 0.001$ ), and showed much less height reduction compared with non-salt treated plants (Fig 5.4f; significant at  $P \leq 0.05$ ).


The number of seeds per plant was not significantly (at  $P \leq 0.05$ ) different between OE lines and WT neither under control (Fig. 5.5a) nor saline (Fig. 5.5b). However, the OE lines showed significant ( $P \leq 0.05$ ) lower reduction in seed number when grown in 300mM NaCl (Fig 5.5c). While the grain yield per plant (g per plant) was not significantly (at  $P \leq 0.05$ ) different between the OE lines and WT under 0 mM conditions (Fig. 5.5d) there was significantly higher grain yield in two of three OE lines under 300 mM (OE-1 and OE-2; significant at  $p \leq 0.001$  and  $p \leq 0.01$  respectively; Fig. 5.5e). This result in a greater maintenance of grain yield in the transgenic lines compared to the control (Fig 5.5f).




**Figure 5.2** Seed size of transgenic lines. The difference in seed size is illustrated for one (OE3 line) of four *35S:AtVHA-C* overexpressing lines under no salt (a, c) and salt stress (b,d) conditions. The other lines show similar results.

**Table 5.1** Seed weight of *AtVHA-C* expressing barley lines and wild type under 0 mM NaCl and 300 mM NaCl growth condition. \*\*\* Significant at  $P \leq 0.001$ . Results are the mean  $\pm$  s.e. (n = 9)


Cultivars	Seed Weight (mg)
WT control	$68.1 \pm 4.0$
OE control	$58.0 \pm 3.0$
WT salt	$15.0 \pm 0.9$
OE salt	$48.1 \pm 3.6^{***}$



**Figure 5.3** Phenotypic difference between non-transformed and overexpressing lines. For simplicity, only two (of four) OE lines are shown. (a-c) Plants treated with 300 mM NaCl for five weeks; (d-f) zero NaCl treatment (no salt stress). The photos were taken 8 weeks after sowing.



**Figure 5.4** Comparative analysis of agronomic characteristics between non-transformed and OE plants. Shoot dry weight (a-c) and plant heights (d-f) are shown for plants grown under zero mM NaCl (no salt; (a, d)) and 300 mM NaCl (saline; b, e) conditions. Panels (c, f) report relative shoot dry weight and plant height, respectively (calculated for salt treated plants as % of control). Results are mean  $\pm$  SE (n=6-15). Measurements are taken 12 weeks after sowing. Asterisks indicate the significant levels of the difference compared with wild type at \*  $P \leq 0.05$ ; \*\*  $P \leq 0.01$ ; and \*\*\*  $P \leq 0.001$ .



**Figure 5.5** Seed number (a-c) and total grain yield (d-f) of OE lines as compared with non-transformed controls. (a, d) – 0 mM NaCl treatment; (b, e) – 300 mM NaCl treatment; (c, f) - relative values (calculated for salt-treated plants as % of control). Results are mean  $\pm$  SE (n=6 to 15). Measurements are taken 12 weeks after sowing. Asterisks indicate the significant levels of the difference compared with wild type at \*  $P \leq 0.05$ ; \*\*  $P \leq 0.01$ ; and \*\*\*  $P \leq 0.001$ .

There was no significant difference in the number of chlorotic leaves under salinity stress between salt-grown OE lines and WT (Fig. 5.6a) but all OE lines showed a significantly ( $P \leq 0.001$ ) lower number of necrotic leaves compared with WT (Fig. 5.6b).


The stomatal conductance of OE lines ranged between 56 and 95  $\text{mmolm}^{-2}\text{s}^{-1}$  with the stomatal conductance of the OE lines significantly lower than the wild type in 0 mM NaCl (Fig. 5.7a). When the plants treated with salt (300 mM NaCl), all three OE lines had approximately 3-fold higher stomatal conductance than WT (17 to 19  $\text{mmolm}^{-2}\text{s}^{-1}$  vs only  $6.1 \pm 0.12$   $\text{mmolm}^{-2}\text{s}^{-1}$  for WT; significant at  $P \leq 0.001$ ; Fig. 5.7b). The chlorophyll content was not significantly different between OE lines and WT under control condition (Fig. 5.7d) but was reduced to a less extent under saline conditions (significant at  $P \leq 0.05$ ; Fig. 5.7f).

### **5.3.3 Transgenic barley expressing *V-ATPase subunit C* has increased leaf $\text{Na}^+$ and $\text{K}^+$ and these ions contributed towards osmotic adjustment**


The OE lines had significantly higher leaf  $\text{Na}^+$  content than WT plants, both under 0 mM NaCl (Fig. 8a) and 300 mM NaCl (Fig. 5.8b) treatments ( $P \leq 0.001$ ). However, the relative increase in leaf  $\text{Na}^+$  under salt stress was less in OE lines compared to WT (six to 7 fold vs 9.2 fold in WT; significant at  $P \leq 0.001$ ; Fig. 5.8c). Leaf  $\text{K}^+$  content for all the OE lines and WT was not significantly (at  $P \leq 0.05$ ) different under control conditions (Fig. 5.8d) but all the OE lines showed much higher leaf  $\text{K}^+$  content compared with WT under saline conditions (significant at  $P \leq 0.001$ ; Fig. 5.8e). Two out of three OE lines (OE-1 and OE-2) have significantly increased the overall leaf  $\text{K}^+$  content (1.5 and 1.4 folds, respectively) while WT plants were not able to do this when grown under saline conditions (Fig. 5.8f).

Osmolality of leaf sap shows there are no differences between OE lines and WT (Table 5.1) but the measured contribution of inorganic ions in WT plants was lower than in OE lines. The relative contribution of inorganic and organic osmolytes towards osmotic adjustment in barley leaves were estimated from these results. In WT, the overall contribution of three

major inorganic osmolytes ( $\text{Na}^+$ ,  $\text{Cl}^-$ , and  $\text{K}^+$ ) was only ~34%, implying that the major bulk of osmotic adjustment under saline conditions (~66%, or two thirds) was achieved by other means, such as *de novo* synthesis of organic osmolytes (Table 5.1). In a stark contrast, contribution of organic osmolytes towards osmotic adjustment in OE lines varied between 49 and 52% (Table 5.1; significant compared with WT a  $P \leq 0.01$ ), e.g. was by ~15% less than in WT.




**Figure 5.6** Difference in the number of chlorotic (a) and necrotic (b) leaves between OE lines and non-transformed barley grown in 300 mM NaCl. Results are mean  $\pm$  SE ( $n=6$  to 15). Measurement was taken 8 weeks after sowing. Asterisks indicate the significant levels of the difference compared with wild type at \*\*\*  $P \leq 0.001$ .



**Figure 5.7** Stomatal conductance (a-c) and leaf chlorophyll content (SPAD values; (d-f)) of OE lines as compared with non-transformed barley. (a, d) – 0 mM NaCl treatment; (b, e) – 300 mM NaCl treatment; (c, f) - relative values (calculated for salt-treated plants as % of control). Results are mean  $\pm$  SE (n=6 to 15). Measurement was taken 8 weeks after sowing. Asterisks indicate the significant levels of the difference compared with wild type at \*  $P \leq 0.05$ ; \*\*  $P \leq 0.01$ ; and \*\*\*  $P \leq 0.001$ .





**Figure 5.8** Leaf Na<sup>+</sup> (a-c) and K<sup>+</sup> (d-f) content of OE lines as compared with non-transformed barley. (a, d) – 0 mM NaCl treatment; (b, e) – 300 mM NaCl treatment; (c, f) - relative values (fold changes in salt-treated plants compared with control). Results are mean ± SE (n=6 to 15). Measurement was taken 8 weeks after sowing. Asterisks indicate the significant levels of the difference compared with wild type at \* P ≤ 0.05; \*\* P ≤ 0.01; and \*\*\* P ≤ 0.001.

**Table 5.2** Relative contribution of Na<sup>+</sup>, K<sup>+</sup> and Cl<sup>-</sup> towards total shoot sap osmolality in transgenic barley leaves grown under saline growth condition. Barley lines designated as *35S:AtVHA-C-1* (OE-1), *35S:AtVHA-C-2* (OE-2), *35S:AtVHA-C-3* (OE-3) and non-transformed control (WT). \* Significant compared with WT at  $P \leq 0.01$ .

	Measured		Calculated	Relative Contribution	
Lines	Total osmolality, mOsm	Contribution of inorganic ions, mOsm	Contribution of organic ions, mOsm	Organic, %	Inorganic, %
OE1	967	492	475.1	49.1	50.9**
OE2	1002	474	527.5	52.6	47.4**
OE3	986	500	485.7	49.3	50.7**
WT	1129	392	737.1	65.3	34.7

## 5.4 Discussion

### 5.4.1 Transgenic *VHA-C* barley improved growth and grain yield under saline growth condition

The activity of V-ATPase increased under salt stress (Silva *et al.* 2010), and both V-ATPase and *NHX* transcripts were up-regulated under salt stress (Qiu *et al.* 2007). Transgenic plants expressing subunits of V-ATPase have shown improvement in growth and better performance under saline condition in several species, such as the c1 subunit in rice (Baisakh *et al.* 2012) and tobacco (Xu *et al.* 2011), as well as a variety of V-ATPase subunits in Arabidopsis (He *et al.* 2014; Wang *et al.* 2011; Zhao *et al.* 2009). All the above examples were related to crops species classified as “salt excluders” which do not normally rely on vacuolar Na<sup>+</sup> sequestration mechanisms (Shabala 2013). For understanding of the growth and physiological response of the *AtVHA-C* gene in barley plants, the gene (*Atlg12840*) was successfully cloned from Arabidopsis cDNA into a transformation vector, under the control of a the *CaMV 35S* promoter and transformed into barley using Agrobacterium mediated transformation. In this

study we show, for the first time, that expressing *VHA-C* from *Arabidopsis* also enhances salinity stress tolerance in a crop plant (barley) which has the ability to accumulate significant levels of shoot  $\text{Na}^+$ .

The transgenic lines expressing *VHA-C* not only showed significant ( $P < 0.05$  and higher) dry weight and plant height when grown under saline conditions (Fig. 5.4) but also higher absolute grain yield (Fig 5.5e). As far as we are aware this is the first work showing transgenic plants expressing V-ATPase subunits have improvements in this important agronomical trait. The closest report we are aware of is improved yield in field grown barley plants expressing the  $\text{H}^+$  V-PPase gene *AVP1* (Schilling *et al.* 2014).

#### **5.4.2 Better stomata control contributes to enhanced salinity tolerance in transgenic plants**

Previous reports suggest the beneficial effects of overexpressing V- $\text{H}^+$ -ATPase transporters (subunits) were related to either improved plant water status, resulting from reduced stomata density and early stomata closure (Baisakh *et al.* 2012), or from increased activity of enzymatic antioxidants (AO) in transgenic lines (He *et al.* 2014). However, plant biomass (and, hence, dry weight) is directly proportional to its ability to assimilate  $\text{CO}_2$  (which relies on gas exchange through open stomates) therefore reductions in stomata density and closure, as suggested by Baisakh *et al.* (2012), cannot be the sole reason *V-H<sup>+</sup>-ATPase* expressing plants have greater salinity tolerance. In addition, there are many reports showing there is no direct correlation between AO activity and salinity stress tolerance in crops (Maksimovic *et al.* 2013). Many salt tolerant plants (such as halophytes) do not require high AO activity as they prevent formation of high levels of reactive oxygen species in the first place (Bose *et al.* 2014a). Therefore must be another mechanism(s) by which *V-H<sup>+</sup>-ATPase* can improve salt tolerance in transgenic plants. From the data reported here, two physiological mechanisms were identified which would lead to improved performance of overexpression lines. The first

one was reduced number of necrotic leaves (Fig 6b) and the second one – higher relative stomatal conductance (Fig 5.7c). The association between coexpressing V-ATPase subunits and stomatal opening and closing has been reported by (Allen *et al.* 2000) using *VHA-C* mutant (*det3*), and also by RNAi knockdown of *OsVHA-A* (Zhang *et al.* 2013). Two to three fold higher Gs values in transgenic lines compared with WT (Fig 5.7b) might account for better CO<sub>2</sub> assimilation, and fewer necrotic leaves increased the overall photosynthetic leaf surface in transgenic place. Both these factors could contribute to increase plant biomass and, ultimately, higher grain yield in OE lines.

#### 5.4.3 Transgenic lines expressing *VHA-C* have better Na<sup>+</sup> sequestration

All the transgenic plants expressing *VHA-C* showed higher leaf Na<sup>+</sup> content than the WT when grown under saline condition (Fig. 5.8b). At the same time, the chlorophyll content was not affected (Fig 5.7), and the number of necrotic leaves was reduced in OE lines (Fig 5.6). Taken together, these facts point out at highly efficient vacuolar Na<sup>+</sup> sequestration in transgenic plants. Na<sup>+</sup>/H<sup>+</sup> antiporters, such as members of the NHX family (Apse *et al.* 2007; Barkla *et al.* 1995; Blumwald *et al.* 1985; Gaxiola *et al.* 1999) energized by the proton motive force generated by proton pumps. The improvement in salinity tolerance in the *V-ATPase C* expressing barley may be due to an improvement in the H<sup>+</sup> gradient between the vacuole and the cytosol, so that native NHX transporters could compartmentalize more Na<sup>+</sup>. Expressing the *subunit C* of the *V-ATPase*, therefore, could be instrumental to increase H<sup>+</sup>-ATPase phosphorylation (Armbruster *et al.* 2004) or stability of its operation (Sze *et al.* 2002), therefore better energizing NHX activity and thus assisting Na<sup>+</sup> sequestration process. Consistent with this model, transgenic Arabidopsis expressing wheat *V-ATPase subunit E* showed increased Na<sup>+</sup> accumulation after 10 days of 120mM NaCl but with a lower Na<sup>+</sup> accumulation in the cytosol (Zhao *et al.* 2009). It remains to be seen whether the improved sequestration in the vacuole is due solely to an increase in the activity of Na<sup>+</sup>/H<sup>+</sup> antiporter

which are already present in the cell or if there is also an up regulation in genes encoding  $\text{Na}^+/\text{H}^+$  antiporters, resulting in a greater concentration (and more activity) of the transporters.

#### 5.4.4 Plants expressing *VHA-C* showed higher $\text{K}^+$ retention

The barley plants expressing *VHA-C* showed significantly higher leaf  $\text{K}^+$  content compared with WT (Fig. 5.8). This feature correlated with the fewer number of necrotic leaves in salt stressed plants (Fig 5.6). This suggests that  $\text{K}^+$  retention in the shoot was essential for preventing salt stress-induced senescence, a programmed cell death process (PCD). The loss of cytosolic  $\text{K}^+$  in cells under stress induces the activation of caspase-like proteases and endonucleases and, hence causes PCD in plants (Demidchik *et al.* 2010; Shabala *et al.* 2007) and mammalian cells (Hughes *et al.* 1998, 1999). Potassium retention in both shoots and roots are crucial mechanisms in salt tolerance in barley which was corroborated by  $\text{K}^+$  kinetics study on shoots and roots of barley and on roots using electrophysiological studies (Adem *et al.* 2014). Recently, Wu *et al.*, (2013) and Wu *et al.*, (2014) showed  $\text{K}^+$  retention in the leaf mesophyll is a contributing factor towards salt tolerance in barley, as measured by non-invasive ion flux. Importantly, a number of  $\text{NHX}$  transporters have been shown to have a higher affinity for  $\text{K}^+$  than  $\text{Na}^+$  (Barragan *et al.* 2012; Bassil *et al.* 2011b; Leidi *et al.* 2010; Rodriguez-Rosales *et al.* 2008) – enhanced proton pumping in *V-H<sup>+</sup>ATPase C* expressing barley may therefore may enable higher accumulation of  $\text{K}^+$  in cell vacuoles. This may contribute to increased cell turgor and better osmotic adjustment under saline conditions (as discussed below). Higher  $\text{K}^+$  content in leaves may be also essential for more efficient stomata control, given the critical role of this nutrient it stomata movements (Anschütz *et al.* 2014).

#### 5.4.5 Relative contribution of organic and inorganic osmolytes

Synthesis of organic osmolytes is energy expensive, particularly in stressed environments (Raven 1985), and plants with enhanced salt tolerance tend to use  $\text{Na}^+$  as cheap osmoticum

for osmotic adjustment (Chen *et al.* 2007a). As shown in Table 5.1, WT plants relied predominantly (65.3% of the cell's osmotic potential) on organic osmolytes in its osmotic adjustment. *V-ATPase C* expressing lines, however, had a large proportion of inorganic osmolytes (specifically,  $\text{Na}^+$  and  $\text{K}^+$ ) to achieve ~ 50% of cell osmotic potential. The transgenic lines were therefore likely to expend less energy generating organic osmolytes and make better use of photosynthetic assimilate (Table 5.1). Osmotic adjustment using inorganic ions is predominant mechanisms in halophytes (Inan *et al.* 2004), and it appears here that some “halophytism” was highly beneficial for transgenic barley plants as well in our case.

## Chapter 6                      General Discussion

Soil salinity is taking nearly three hectares of arable land from conventional crop farming every minute (Shabala *et al.* 2014). Of the world's 5.2 billion ha land coverage, 3.6 billion is dryland. Dryland is impacted by erosion, soil degradation and salinization (Riadh *et al.* 2010). Half of the world's irrigated land is compromised by salinization, alkalization or waterlogging (Szabolcs 1994). In Australia, the clearing of vegetation for farmland that started a century ago has contributed to saline soil covering 2 million ha, with estimates that another 15 million ha are at risk of turning saline in the next 50 years (National Land and Water Resources Audit; <http://audit.ea.gov.au>). It is estimated that currently one-third of Australian soil and 67% of Australia's agricultural land has the potential for transient salinity (Rengasamy 2006).

The salinity problem can be tackled in a number of ways. One could involve changing farm management practices, such as efficient use of irrigation. For example, drip irrigation conserves water-use and hence reduces salinization. Meanwhile in rain-fed agriculture, rotation of annual crops with deep rooted perennial plants prevents the water table from rising thus reducing transfer of the salt to the surface. Salt tolerant plants are the most cost effective solution in combating salinization (Munns 2002). In broader term, the solution to the problem is two-fold. The first option is to intensify crop production using currently available arable land by improving the productivity of crops exposed to stress. This calls for a major leap in breeding crops for stress tolerance. This option is limited by the availability of genetic diversity, in other words, a smaller gene pool to be tapped into to develop stress tolerant crops (Colmer *et al.* 2005). The second option could be identification of genes from extremophiles and introducing these into traditional crops (Shabala 2013; Shabala *et al.* 2014).

Salinity tolerance in crops is a multifaceted physiological trait involving multiple mechanisms. The main ones are osmotic adjustment, minimizing Na<sup>+</sup> uptake by roots and/or increasing Na<sup>+</sup> efflux back to the soil, intracellular Na<sup>+</sup> sequestration, potassium retention in

the cytosol, tissue-specific  $\text{Na}^+$  sequestration, control of xylem ion loading, excluding  $\text{Na}^+$  from the shoot and oxidative stress tolerance (Munns *et al.* 2008; Shabala *et al.* 2012; Zhu 2003). This list of mechanisms can be grouped into three major clusters: (i) osmo-tolerance; (ii) sodium exclusion; and (iii) tissue tolerance mechanisms (Munns *et al.* 2008). In spite of the fact that significant progress has been made in explaining specific details of each of these mechanisms, the relative contribution of the afore-mentioned components to overall salinity tolerance remains unclear. Part of this thesis study, efforts were made to elucidate relative contribution of the mechanisms of salt tolerance in barley, with experiments performed that spanned from the cellular to the whole plant level.

To begin with, crop osmotolerance has been attributed to a plant's ability to increase *de novo* synthesis of compatible solutes (Chen *et al.* 2007a; Hasegawa *et al.* 2000; Munns *et al.* 2015; Shabala *et al.* 2011b). This has been a less efficient way towards salt tolerance, as it involves energy expensive ATP to synthesise osmolytes (Chen *et al.* 2007a). At least for salt tolerant crops like barley, the usage of osmolytes for osmotic adjustment is less likely to be the primary mechanism due to the energy cost instead  $\text{Na}^+$  is preferentially used as a “cheap” osmoticum. The other major component of salinity tolerance is  $\text{Na}^+$  exclusion. This mechanism employs the plasma membrane  $\text{Na}^+/\text{H}^+$  exchanger SOS1, which is activated through a  $\text{Ca}^{2+}$  dependent signaling pathway via phosphorylation by interacting proteins SOS2 and SOS3 (Shi *et al.* 2003; Wu *et al.* 1996). This process energized by the plasma membrane  $\text{H}^+$ -ATPase (Palmgren *et al.* 2011). The  $\text{Na}^+$  extruded to the medium by this mechanism however can enhance osmotic and ionic imbalance that exacerbates the existing salinity stress. This mechanism can only be used as a short-term solution and may not confer salinity tolerance under field conditions. The other mechanisms restrict  $\text{Na}^+$  accumulation in shoots involve reducing  $\text{Na}^+$  loading into the xylem (Munns *et al.* 2008; Tester *et al.* 2003) and increasing  $\text{Na}^+$  retrieval from the xylem (Davenport *et al.* 2007). Recently, Munns *et al.*,



(2012) reported 25% grain yield increases in durum wheat due to the presence of *TmHKT1;5*-A introgressed into commercial durum wheat cultivar. The gene is responsible for retrieving  $\text{Na}^+$  from the xylem. The other mechanism involves sodium retrieval from the shoot via recirculation to the root via phloem (Berthomieu *et al.* 2003). This mechanism is believed to use the sodium HKT transporter (Garcia-deblas *et al.* 2003). Nevertheless, it's debated that excluding  $\text{Na}^+$  from the xylem may not always be a plausible mechanism (Shabala *et al.* 2010). It's because some species that employ high tissue tolerance mechanism use  $\text{Na}^+$  as a cheap osmoticum. In light of this, *HvHKT2;1* over-expressing barley plants showed higher xylem and leaf  $\text{Na}^+$  content in saline conditions and showed increased salt tolerance (Mian *et al.* 2011). Our findings also provide a physiological evidence that higher shoot sodium correlates with tolerance in barley (Adem *et al.* 2014). This is explained as the ability of the plant to accumulate large amounts of sodium in the tissue without causing harm to cellular functions. This could be achieved by sequestering excess  $\text{Na}^+$  away from the cytosol into the vacuole or away from the actively photosynthesizing tissue. This is effected by the tonoplast  $\text{Na}^+/\text{H}^+$  exchanger (Apse *et al.* 1999; Blumwald 2000; Gaxiola *et al.* 1999) energized by vacuolar  $\text{H}^+$ -ATPase and vacuolar pyrophosphatase (Fukuda *et al.* 2004a).

Tissue tolerance is one of the various mechanisms used by plants for salt tolerance and is a major contributor to salt tolerance in barley. This tolerance mechanism involves sodium sequestration away from the cytosol where sodium is deleterious for normal cellular metabolism. Sodium ions are the main contributor to the disturbance of cell metabolism, enzymatic activity and ionic homeostasis (Ahmad *et al.* 2014; Anschütz *et al.* 2014; Shabala *et al.* 2014). For a plant to grow under saline conditions, sodium should be excluded from the shoot keeping the  $\text{K}^+$  at high levels thereby maintaining a high  $\text{K}^+/\text{Na}^+$  ratio, particularly in the leaves (Ren *et al.* 2005; Serrano *et al.* 2001; Shi *et al.* 2002a). To achieve the maintenance of high  $\text{K}^+/\text{Na}^+$  ratios in the cytosol, plants extrude  $\text{Na}^+$  out of the cell via the plasma

membrane and/or sequestering  $\text{Na}^+$  in vacuoles. The compartmentalization process is vital for preventing  $\text{Na}^+$  toxicity in the cytosol and for osmotic adjustment of a cell conferring salt tolerance (Blumwald *et al.* 2000).  $\text{Na}^+$  is transported across the tonoplast into the vacuole by exchanging protons. A gene encoding the  $\text{Na}^+/\text{H}^+$  exchanger *NHX1* was found to compartmentalize  $\text{Na}^+$  in prevacuolar and vacuolar compartments (Nass *et al.* 1997; Nass *et al.* 1998). The NHX protein is located in the tonoplast and energized by Vacuolar  $\text{H}^+$ -ATPase and Vacuolar  $\text{H}^+$ -inorganic pyrophosphatase (Gaxiola *et al.* 2007; Sze *et al.* 1999). Expressing *AtNHX1* gene was reported to confer salt tolerance in salt sensitive categories of plants (Apse *et al.* 1999; Bayat *et al.* 2011; Wu *et al.* 2009; Wu *et al.* 2004; Xue *et al.* 2004; Zhang *et al.* 2001a; Zhang *et al.* 2001b). Recently, a new role has emerged for the NHX1 protein in that it also transports  $\text{K}^+$  ions (Barragan *et al.* 2012; Bassil *et al.* 2011b). Despite the above literature claims, in this PhD study expressing *AtNHX1* gene in barley did not confer any additional salt tolerance. There are several possible explanations for why this is the case. One reason could be a misfolding of *AtNHX1* protein rendering it inactive. The second reason could be due to mistargeting of the protein into other membranes. A similar phenomenon was observed in the over-expression of *AVP1* which was mistargeted to the plasma membrane (Gaxiola *et al.* 2012; Li *et al.* 2005). The third reason might involve lack of required post-translational modification such as phosphorylation. Kinases, such as SOS2, are important regulatory components in other salt tolerance signaling systems (Neuhaus *et al.* 2014; Weinl *et al.* 2009). A more physiological oriented reason may comprise back leak of  $\text{Na}^+$  from the vacuole to the cytosol via  $\text{Na}^+$  permeable SV (slow activating) and FV (fast activating) channels, which typically confer salt tolerance in halophyte plant (Bonales-Alatorre *et al.* 2013b; Hedrich *et al.* 1987; Shabala 2013). Another physiological reason is the importance on needing to have highly active V-ATPase (Krebs *et al.* 2010; Vera-Estrella *et al.* 1999; Wang *et al.* 2001) and V-PPiase (Guo *et al.* 2006; Parks *et al.* 2002; Vera-Estrella *et al.* 2005) necessary to support the

energy needs of the  $\text{Na}^+$  sequestration process by NHX ((Shabala 2013). In the transgenic barley plants activities of these enzymes may have been insufficient to support enhanced  $\text{Na}^+$  sequestration (Adem *et al.* 2015). Furthermore, a reduced  $\text{K}^+$  pool in the cytosol resulting from its enhanced sequestration in the vacuole might in turn require osmolyte production to prevent dehydration, a process which is energetically costly and thus would lead to yield penalties. However, the compensation for the cytosolic  $\text{K}^+$  loss by osmolyte production is not the case for tolerant barley genotypes (Chen *et al.* 2007a) and the genetic background of the barley plant used in this study was a comparatively tolerant cultivar (Adem *et al.* 2014). Sequestration of  $\text{Na}^+$  in the vacuole is possible by  $\text{Na}^+$  loading, energization of loading and retention (Shabala 2013). In light of this, it was shown that better salt tolerance was obtained by co-expression of the *AVP* with *NHX* (Zhao *et al.* 2006). Hence, studying the individual components for sequestration and pyramiding them would be highly advantageous. Therefore, studies are showing that over-expression of *AVP1* improve tolerance to salinity and drought in *Arabidopsis* (Gaxiola *et al.* 2002; Gaxiola *et al.* 2001; Gaxiola *et al.* 2012; Guo *et al.* 2006). Similarly, the over-expression of VHA subunits has shown salt tolerant phenotypes. For instance, V-ATPase c1 subunit from halophyte grass improved rice growth under saline conditions (Baisakh *et al.* 2012). He *et al.*, (2014) and Wang *et al.*, (2011) reported the improvement of salt tolerance by cloning wheat V-ATPase subunit B in *Arabidopsis*. Furthermore, Zhao *et al.*, (2009) has shown salt tolerant plants by over-expressing V-ATPase subunit E in *Arabidopsis*. Similarly, salt tolerant plants were obtained by over-expressing V-ATPase subunit c (Xu *et al.* 2011). In our study, we have shown the improvement of biomass and grain yield of barley plants by expressing *Arabidopsis thaliana* V-ATPase subunit C (*AtVHA-C*). The improvement in grain yield and biomass in our V-ATPase transgenic lines was as a result of better osmotic adjustment, as shown by better accumulation of  $\text{Na}^+$  and  $\text{K}^+$  in the shoot and higher contribution of inorganic ions ( $\text{Na}^+$  and  $\text{Cl}^-$ ) to osmotic adjustment

than organic solutes. In this PhD study, it was shown there was a lack of phenotype when *AtNHX1* was expressed on its own whereas a salt tolerance phenotype was observed when *AtVHA-C* was expressed. Given that co-expression of *NHX* gene with genes that encode proton pumps have been shown to have the best phenotypes (Zhao *et al.* 2006), the two transgenic genotypes used in this study could be crossed to pyramid both the *AtNHX1* and *AtVHA-C* and the subsequent offspring tested to see if pyramiding these genes might result in a better salt tolerant barley plant than simply expressing *AtVHA-C* alone. Such a combination may also help to investigate the reason behind why no phenotype was seen in *AtNHX1* expressing plants. It might answer the questions as to whether there was not enough protons being pumped into the vacuole in the *NHX1* expressing barley for the *NHX* to work. The *AtVHA-C* and the combined genotype expressing both *AtNHX1* and *AtVHA-C* need to be tested in the field in multiple sites and for multiple years to see if they show grain yield improvement. The *AtVHA-C* barley plants and the pyramided barley plants containing these two genes could be used for farmers if they show consistent results in future field trials or they can be used to show the importance of these two genes in yield improvement in saline soils and thus representing a stepping stone to more non-transgenic options. This could be achieving by screening for the best barley germplasm that has good expression of the native *VHA-C* gene and develop a marker to the gene so it can be introgressed into an elite barley cultivar using conventional breeding method. An alternative strategy would be to use the CRISPR (clustered regularly interspaced short palindromic repeats)- Cas system which can be used to selectively modify or edits a target region (Cheng *et al.* 2013; Sander *et al.* 2014; Shan *et al.* 2013). In the case here this would involve modifying the promoter region of the *VHA-C* gene to constitutively express the gene without requiring the development of a transgenic barley plant. The CRISPR-Cas system is a recent invention and there is still an ongoing debate about whether crops developed using this system are GMOs or non-GMOs. It

is a potentially superior option to rapidly obtain and transfer plants with desired traits to breeders and producers.

Vacuolar  $\text{Na}^+$  sequestration is not the only mechanism contributing to tissue tolerance. Potassium retention is also emerging as an important tissue tolerance mechanism in root (Chen *et al.* 2005; Chen *et al.* 2007b; Chen *et al.* 2008; Chen *et al.* 2007d) and leave tissues (Wu *et al.* 2013). The maintenance of high  $\text{K}^+$  in the cytosol suppresses the activity of caspase like proteolytic and endonucleolytic enzymes triggering programmed cell death in salt affected cells (Demidchik *et al.* 2010; Shabala *et al.* 2007). Cytosolic  $\text{K}^+$  plays roles in assisting metabolic processes, such as protein synthesis, by enabling the binding of tRNA to ribosomes (Wyn Jones *et al.* 1979).

An additional mechanism that contributes to salinity tolerance under tissue tolerance is oxidative stress tolerance. Studies have shown that significant amounts of ROS are generated in salt affected root and leaf tissues (Miller *et al.* 2008; Mittler 2002). The causal link between salinity and oxidative stress signaling and reactive species detoxification has emerged (Bose *et al.* 2014a). Our findings have shown improved prevention of hydroxyl radical- induced  $\text{K}^+$  loss in tolerant barley cultivars. This can be an indication of cross protection between stress responses in this case oxidative stress with salinity stress. This finding is also consistent with the report on root response to  $\text{H}_2\text{O}_2$  (Maksimovic *et al.* 2013). NADPH oxidase is responsible for the ROS generation under salinity stress (Sagi *et al.* 2006). Accordingly, we studied the barley homologue of *RBoHF* gene transcript level and showed the least down regulation in the leaves of salt sensitive cultivar, which suggests a causal link between NADPH oxidase activity and tissue tolerance. Unlike in the leaves, the root transcript level compared to the control was non-significant (Adem *et al.* 2014). In our study, the transcript level of barley homologue *NHX*, *RBoHF*, *AHA* and *GORK* genes that are responsible for  $\text{Na}^+/\text{H}^+$  transport to the vacuole, NADPH oxidase (ROS generation), plasma membrane  $\text{H}^+$  pumping and outward

rectified  $K^+$  efflux, respectively were studied at one point in time at seedling stage in both roots and leaves. At the tested time point and growth stage, the transcript level was found not to be a good indicator of salinity tolerance. This result may be interpreted in a number of ways. First, salt tolerance genes were shown to fluctuate between days (e.g. *HVP10*) (Shavrukov *et al.* 2013). Second, the transcript amount may not always have a predictive value towards salinity tolerance at the whole plant level or a cellular level, since the gene product could be dependent upon post translational modification such as phosphorylation and proper protein folding. Taken together, the above mentioned genes need to be studied at different time scale and in different growth stage of the shoot and in different region of the root to better understand their roles in plant salinity tolerance.

To recap, soil salinity is a major abiotic stress costing billions of dollars to the agricultural industry and to combat this stress, a number of strategies can be devised, however, the cost effective method is to improve crop salt tolerance through genetic manipulation. Improvement of salt tolerance in barley should focus on tissue tolerance mechanisms. Sodium sequestration is one of the tissue tolerance mechanism and this study has indicated that different approaches must be used in improving salt tolerance where the *NHX* expression shown to not benefiting barley plants in improving salt tolerance unlike salt excluder crops. In light of this, the future efforts in sodium sequestration especially in a salt including crop like barley ought to consider co-expression of other genes involved in the sequestration process. Results have also shown that genes with specific function like *NHX* need the help of other genes such as pumps and those involved in  $Na^+$  leaks. However, improvements in halotolerance using genes with general functions, for instance, proton pumps, as shown by expressing *AtVHA-C* (our study) and *AVP* in barley (Schilling *et al.* 2014), has shown to improve salt tolerance when expressed as single gene. Hence, the nature of the gene to be

---

expressed must be looked into first before we express it as single gene or co-express it with other genes.

## References

- Adem, G, Roy, S, Plett, D, Zhou, M, Bowman, J & Shabala, S 2015, 'Expressing *AtNHX1* in barley (*Hordium vulgare* L.) does not improve plant performance under saline conditions', *Plant Growth Regulation*, pp. 1-9.
- Adem, GD, Roy, SJ, Zhou, M, Bowman, JP & Shabala, S 2014, 'Evaluating contribution of ionic, osmotic and oxidative stress components towards salinity tolerance in barley', *BMC Plant Biology*, vol. 14.
- Adler, G, Blumwald, E & Bar-Zvi, D 2010, 'The sugar beet gene encoding the sodium/proton exchanger 1 (*BvNHX1*) is regulated by a MYB transcription factor', *Planta*, vol. 232, no. 1, pp. 187-195.
- Adolf, VI, Shabala, S, Andersen, MN, Razzaghi, F & Jacobsen, SE 2012, 'Varietal differences of quinoa's tolerance to saline conditions', *Plant and Soil*, vol. 357, no. 1-2, pp. 117-129.
- Aharon, GS, Apse, MP, Duan, SL, Hua, XJ & Blumwald, E 2003, 'Characterization of a family of vacuolar  $\text{Na}^+/\text{H}^+$  antiporters in *Arabidopsis thaliana*', *Plant and Soil*, vol. 253, no. 1, pp. 245-256.
- Ahmad, I & Maathuis, FJM 2014, 'Cellular and tissue distribution of potassium: Physiological relevance, mechanisms and regulation', *Journal of Plant Physiology*, vol. 171, no. 9, pp. 708–714.
- Al-Naseri, A, Bowman, JP, Wilson, R, Nilsson, RE & Britz, ML 2013, 'Impact of Lactose Starvation on the Physiology of *Lactobacillus casei* GCRL163 in the Presence or Absence of Tween 80', *Journal of Proteome Research*.
- Allen, GJ, Chu, SP, Schumacher, K, Shimazaki, CT, Vafeados, D, Kemper, A, Hawke, SD, Tallman, G, Tsien, RY, Harper, JF, Chory, J & Schroeder, JI 2000, 'Alteration of stimulus-specific guard cell calcium oscillations and stomatal closing in *Arabidopsis det3* mutant', *Science*, vol. 289, no. 5488, pp. 2338-2342.
- Anschütz, U, Beckera, D & Shabala, S 2014, 'Going beyond nutrition: Regulation of potassium homeostasis as a common denominator of plant adaptive responses to environment', *Journal of Plant Physiology*, vol. 171, pp. 670–687.
- Apse, MP, Aharon, GS, Snedden, WA & Blumwald, E 1999, 'Salt tolerance conferred by overexpression of a vacuolar  $\text{Na}^+/\text{H}^+$  antiport in *Arabidopsis*', *Science*, vol. 285, no. 5431, pp. 1256-1258.
- Apse, MP & Blumwald, E 2007, ' $\text{Na}^+$  transport in plants', *FEBS Letters*, vol. 581, no. 12, pp. 2247-2254.
- Apse, MP, Sottosanto, JB & Blumwald, E 2003, 'Vacuolar cation/ $\text{H}^+$  exchange, ion homeostasis, and leaf development are altered in a T-DNA insertional mutant of *AtNHX1*, the *Arabidopsis* vacuolar  $\text{Na}^+/\text{H}^+$  antiporter', *Plant Journal*, vol. 36, no. 2, pp. 229-239.



- Armbruster, A, Svergun, DI, Coskun, U, Juliano, S, Bailer, SM & Gruber, G 2004, 'Structural analysis of the stalk subunit Vma5p of the yeast V-ATPase in solution', *FEBS Letters*, vol. 570, no. 1-3, pp. 119-125.
- Ayala-Navarrete, LI, Mechanicos, AA, Gibson, JM, Singh, D, Bariana, HS, Fletcher, J, Shorter, S & Larkin, PJ 2013, 'The Pontin series of recombinant alien translocations in bread wheat: single translocations integrating combinations of *Bdv2*, *Lr19* and *Sr25* disease-resistance genes from *Thinopyrum intermedium* and *Th. ponticum*', *Theoretical and Applied Genetics*, vol. 126, no. 10, pp. 2467-2475.
- Baisakh, N, RamanaRao, MV, Rajasekaran, K, Subudhi, P, Janda, J, Galbraith, D, Vanier, C & Pereira, A 2012, 'Enhanced salt stress tolerance of rice plants expressing a vacuolar H<sup>+</sup>-ATPase subunit c1 (*SaVHAc1*) gene from the halophyte grass *Spartina alterniflora* Loisel', *Plant Biotechnology Journal*, vol. 10, no. 4, pp. 453-464.
- Banks, PM, Larkin, PJ, Bariana, HS, Lagudah, ES, Appels, R, Waterhouse, PM, Brettell, RIS, Chen, X, Xu, HJ, Xin, ZY, Qian, YT, Zhou, XM, Cheng, ZM & Zhou, GH 1995, 'The use of cell-culture for subchromosomal introgressions of barley yellow dwarf virus resistance from *Thinopyrum intermedium* to wheat', *Genome*, vol. 38, no. 2, pp. 395-405.
- Barkla, BJ, Vera-Estrella, R, Hernandez-Coronado, M & Pantoja, O 2009, 'Quantitative proteomics of the tonoplast reveals a role for glycolytic enzymes in salt tolerance', *Plant Cell*, vol. 21, no. 12, pp. 4044-4058.
- Barkla, BJ, Zingarelli, L, Blumwald, E & Smith, JAC 1995, 'Tonoplast Na<sup>+</sup>/H<sup>+</sup> antiport activity and its energization by the vacuolar H<sup>+</sup>-ATPase in the halophytic plant *Mesembryanthemum crystallinum* L.', *Plant Physiology*, vol. 109, no. 2, pp. 549-556.
- Barragan, V, Leidi, EO, Andres, Z, Rubio, L, De Luca, A, Fernandez, JA, Cubero, B & Pardo, JM 2012, 'Ion exchangers NHX1 and NHX2 Mediate active potassium uptake into vacuoles to regulate cell turgor and stomatal function in Arabidopsis', *Plant Cell*, vol. 24, no. 3, pp. 1127-1142.
- Barrett-Lennard, E & Setter, T 2010, 'Developing saline agriculture: from traits and genes to systems', *Functional Plant Biology*, vol. 37, pp. iii-iv.
- Bassil, E, Ohto, MA, Esumi, T, Tajima, H, Zhu, Z, Cagnac, O, Belmonte, M, Peleg, Z, Yamaguchi, T & Blumwald, E 2011a, 'The Arabidopsis intracellular Na<sup>+</sup>/H<sup>+</sup> antiporters NHX5 and NHX6 are endosome associated and necessary for plant growth and development', *Plant Cell*, vol. 23, no. 1, pp. 224-239.
- Bassil, E, Tajima, H, Liang, YC, Ohto, MA, Ushijima, K, Nakano, R, Esumi, T, Coku, A, Belmonte, M & Blumwald, E 2011b, 'The Arabidopsis Na<sup>+</sup>/H<sup>+</sup> antiporters NHX1 and NHX2 control vacuolar pH and K<sup>+</sup> homeostasis to regulate growth, flower development, and reproduction', *Plant Cell*, vol. 23, no. 12, pp. 3482-3497.
- Batelli, G, Verslues, PE, Agius, F, Qiu, Q, Fujii, H, Pan, S, Schumaker, KS, Grillo, S & Zhu, JK 2007, '*SOS2* promotes salt tolerance in part by interacting with the vacuolar H<sup>+</sup>-ATPase and upregulating its transport activity', *Molecular and Cellular Biology*, vol. 27, no. 22, pp. 7781-7790.

- Bayat, F, Shiran, B & Belyaev, DV 2011, 'Overexpression of *HvNHX2*, a vacuolar  $\text{Na}^+/\text{H}^+$  antiporter gene from barley, improves salt tolerance in *Arabidopsis thaliana*', *Australian Journal of Crop Science*, vol. 5, pp. 428-432.
- Berthomieu, P, Conejero, G, Nublat, A, Brackenbury, WJ, Lambert, C, Savio, C, Uozumi, N, Oiki, S, Yamada, K, Cellier, F, Gosti, F, Simonneau, T, Essah, PA, Tester, M, Very, AA, Sentenac, H & Casse, F 2003, 'Functional analysis of *AtHKT1* in *Arabidopsis* shows that  $\text{Na}^+$  recirculation by the phloem is crucial for salt tolerance', *Embo Journal*, vol. 22, no. 9, pp. 2004-2014.
- Bian, M, Waters, I, Broughton, S, Zhang, XQ, Zhou, MX, Lance, R, Sun, DF & Li, CD 2013, 'Development of gene-specific markers for acid soil/aluminium tolerance in barley (*Hordeum vulgare* L.)', *Molecular Breeding*, vol. 32, no. 1, pp. 155-164.
- Blumwald, E 2000, 'Sodium transport and salt tolerance in plants', *Current Opinion in Cell Biology*, vol. 12, no. 4, pp. 431-434.
- Blumwald, E, Aharon, GS & Apse, MP 2000, 'Sodium transport in plant cells', *Biochimica Et Biophysica Acta-Biomembranes*, vol. 1465, no. 1-2, pp. 140-151.
- Blumwald, E & Poole, RJ 1985, ' $\text{Na}^+/\text{H}^+$  antiport in isolated tonoplast vesicles from storage tissue of *Beta vulgaris*', *Plant Physiology*, vol. 78, no. 1, pp. 163-167.
- Bolwell, GP & Wojtaszek, P 1997, 'Mechanisms for the generation of reactive oxygen species in plant defence - a broad perspective', *Physiological and Molecular Plant Pathology*, vol. 51, no. 6, pp. 347-366.
- Bonales-Alatorre, E, Pottosin, I, Shabala, L, Chen, Z-H, Zeng, F, Jacobsen, S-E & Shabala, S 2013a, 'Differential activity of plasma and vacuolar membrane transporters contributes to genotypic differences in salinity tolerance in halophyte species, *Chenopodium quinoa*', *International Journal of Molecular Sciences*, vol. 14, pp. 9267-9285.
- Bonales-Alatorre, E, Shabala, S, Chen, Z-H & Pottosin, I 2013b, 'Reduced tonoplast fast-activating and slow-activating channel activity is essential for conferring salinity tolerance in a facultative halophyte, quinoa', *Plant Physiology*, vol. 162, pp. 1-13.
- Bose, J, Rodrigo-Moreno, A & Shabala, S 2014a, 'ROS homeostasis in halophytes in the context salinity stress tolerance', *Journal of Experimental Botany*, vol. 65, no. 5, pp. 1241-1257.
- Bose, J, Shabala, L, Pottosin, I, Zeng, FR, Velarde-Buendia, AM, Massart, A, Poschenrieder, C, Hariadi, Y & Shabala, S 2014b, 'Kinetics of xylem loading, membrane potential maintenance, and sensitivity of  $\text{K}^+$ -permeable channels to reactive oxygen species: physiological traits that differentiate salinity tolerance between pea and barley', *Plant Cell and Environment*, vol. 37, no. 3, pp. 589-600.
- Brown, M & Funk, C 2008, 'Climate-Food Security under climate change', *Science*, vol. 319, pp. 580-581.
- Bruggemann, LI, Pottosin, II & Schonknecht, G 1999, 'Selectivity of the fast activating vacuolar cation channel', *Journal of Experimental Botany*, vol. 50, no. 335, pp. 873-876.

- Brummer, B, Warncke, J & Slayman, C 1985, 'Control of membrane permeability by extracellular calcium pump and leak behavior of the *Neurospora* plasma membrane', *Plant Physiology*, vol. 77, pp. S-143.
- Brüx, A, Liu, TY, Krebs, M, Stierhof, YD, Lohmann, JU, Miersch, O, Wastermack, C & Schumacher, K 2008, 'Reduced V-ATPase activity in the trans-Golgi network causes oxylipin-dependent hypocotyl growth inhibition in *Arabidopsis*', *Plant Cell*, vol. 20, no. 4, pp. 1088-1100.
- Burton, RA, Jobling, SA, Harvey, AJ, Shirley, NJ, Mather, DE, Bacic, A & Fincher, GB 2008, 'The genetics and transcriptional profiles of the cellulose synthase-like *HvCslF* gene family in barley', *Plant Physiology*, vol. 146, no. 4, pp. 1821-1833.
- Carpaneto, A, Cantu, AM & Gambale, F 1999, 'Redox agents regulate ion channel activity in vacuoles from higher plant cells', *FEBS Letters*, vol. 442, no. 2-3, pp. 129-132.
- Chazen, O, Hartung, W & Neumann, PM 1995, 'The different effects of PEG-6000 and NaCl on leaf development are associated with differential inhibition of root water transport', *Plant Cell and Environment*, vol. 18, no. 7, pp. 727-735.
- Chen, Z, Cuin, TA, Zhou, M, Twomey, A, Naidu, BP & Shiabala, S 2007a, 'Compatible solute accumulation and stress-mitigating effects in barley genotypes contrasting in their salt tolerance', *Journal of Experimental Botany*, vol. 58, no. 15-16, pp. 4245-4255.
- Chen, Z, Newman, I, Zhou, M, Mendham, N, Zhang, G & Shabala, S 2005, 'Screening plants for salt tolerance by measuring  $K^+$  flux: a case study for barley', *Plant Cell and Environment*, vol. 28, no. 10, pp. 1230-1246.
- Chen, Z, Pottosin, I, Cuin, TA, Fuglsang, AT, Tester, M, Jha, D, Zepeda-Jazo, I, Zhou, MX, Palmgren, MG, Newman, IA & Shabala, S 2007b, 'Root plasma membrane transporters controlling  $K^+/Na^+$  homeostasis in salt-stressed barley', *Plant Physiology*, vol. 145, no. 4, pp. 1714-1725.
- Chen, Z, Shabala, S, Mendham, N, Newman, I, Zhang, G & Zhou, M 2008, 'Combining ability of salinity tolerance on the basis of NaCl-induced  $K^+$  flux from roots of barley', *Crop Science*, vol. 48, no. 4, pp. 1382-1388.
- Chen, Z, Zhou, M, Newman, I, Mendham, N, Zhang, G & Shabala, S 2007c, 'Potassium and sodium relations in salinised barley tissues as a basis of differential salt tolerance', *Functional Plant Biology*, vol. 34, no. 2, pp. 150-162.
- Chen, ZH, Zhou, M, Newman, IA, Mendham, NJ, Zhang, G & Shabala, S 2007d, 'Potassium and sodium relations in salinised barley tissues as a basis of differential salt tolerance', *Functional Plant Biology*, vol. 34, no. 2, pp. 150-162.
- Cheng, AW, Wang, HY, Yang, H, Shi, LY, Katz, Y, Theunissen, TW, Rangarajan, S, Shivalila, CS, Dadon, DB & Jaenisch, R 2013, 'Multiplexed activation of endogenous genes by CRISPR-on, an RNA-guided transcriptional activator system', *Cell Research*, vol. 23, no. 10, pp. 1163-1171.
- Chomczynski, P 1993, 'A reagent for the single-step simultaneous isolation of RNA, DNA and Protein from cell and tissue samples', *Biotechniques*, vol. 15, no. 3, pp. 532-537.

- Colmer, TD, Munns, R & Flowers, TJ 2005, 'Improving salt tolerance of wheat and barley: future prospects', *Australian Journal of Experimental Agriculture*, vol. 45, no. 11, pp. 1425-1443.
- Cuin, TA, Bose, J, Stefano, G, Jha, D, Tester, M, Mancuso, S & Shabala, S 2011, 'Assessing the role of root plasma membrane and tonoplast  $\text{Na}^+/\text{H}^+$  exchangers in salinity tolerance in wheat: in planta quantification methods', *Plant Cell and Environment*, vol. 34, no. 6, pp. 947-961.
- Cuin, TA, Parsons, D & Shabala, S 2010, 'Wheat cultivars can be screened for NaCl salinity tolerance by measuring leaf chlorophyll content and shoot sap potassium', *Functional Plant Biology*, vol. 37, no. 7, pp. 656-664.
- Cuin, TA & Shabala, S 2007, 'Compatible solutes reduce ROS-induced potassium efflux in Arabidopsis roots', *Plant Cell and Environment*, vol. 30, no. 7, pp. 875-885.
- Cuin, TA, Tian, Y, Betts, SA, Chalmandrier, R & Shabala, S 2009, 'Ionic relations and osmotic adjustment in durum and bread wheat under saline conditions', *Functional Plant Biology*, vol. 36, no. 12, pp. 1110-1119.
- Curtis, MD & Grossniklaus, U 2003, 'A gateway cloning vector set for high-throughput functional analysis of genes in planta', *Plant Physiology*, vol. 133, no. 2, pp. 462-469.
- Dai, F, Nevo, E, Wu, DZ, Comadran, J, Zhou, MX, Qiu, L, Chen, ZH, Beiles, A, Chen, GX & Zhang, GP 2012, 'Tibet is one of the centers of domestication of cultivated barley', *Proceedings of the National Academy of Sciences of the United States of America*, vol. 109, no. 42, pp. 16969-16973.
- Davenport, R, James, RA, Zakrisson-Plogander, A, Tester, M & Munns, R 2005, 'Control of sodium transport in durum wheat', *Plant Physiology*, vol. 137, no. 3, pp. 807-818.
- Davenport, RJ, Munoz-Mayor, A, Jha, D, Essah, PA, Rus, A & Tester, M 2007, 'The  $\text{Na}^+$  transporter *AtHKT1;1* controls retrieval of  $\text{Na}^+$  from the xylem in Arabidopsis', *Plant Cell and Environment*, vol. 30, no. 4, pp. 497-507.
- Davenport, RJ, Reid, RJ & Smith, FA 1997, 'Sodium-calcium interactions in two wheat species differing in salinity tolerance', *Physiologia Plantarum*, vol. 99, no. 2, pp. 323-327.
- Demidchik, V, Cuin, TA, Svistunenko, D, Smith, SJ, Miller, AJ, Shabala, S, Sokolik, A & Yurin, V 2010, 'Arabidopsis root  $\text{K}^+$ -efflux conductance activated by hydroxyl radicals: single-channel properties, genetic basis and involvement in stress-induced cell death', *Journal of Cell Science*, vol. 123, no. 9, pp. 1468-1479.
- Demidchik, V, Davenport, RJ & Tester, M 2002, 'Nonselective cation channels in plants', *Annual Review of Plant Biology*, vol. 53, pp. 67-107.
- Demidchik, V, Shabala, SN, Coutts, KB, Tester, MA & Davies, JM 2003, 'Free oxygen radicals regulate plasma membrane  $\text{Ca}^{2+}$  and  $\text{K}^+$ -permeable channels in plant root cells', *Journal of Cell Science*, vol. 116, no. 1, pp. 81-88.

- Dettmer, J, Schubert, D, Calvo-Weimar, O, Stierhof, YD, Schmidt, R & Schumacher, K 2005, 'Essential role of the V-ATPase in male gametophyte development', *Plant Journal*, vol. 41, no. 1, pp. 117-124.
- Dietz, KJ, Tavakoli, N, Kluge, C, Mimura, T, Sharma, SS, Harris, GC, Chardonnens, AN & Golldack, D 2001, 'Significance of the V-type ATPase for the adaptation to stressful growth conditions and its regulation on the molecular and biochemical level', *Journal of Experimental Botany*, vol. 52, no. 363, pp. 1969-1980.
- Dobrovinskaya, OR, Muniz, J & Pottosin, II 1999, 'Inhibition of vacuolar ion channels by polyamines', *Journal of Membrane Biology*, vol. 167, no. 2, pp. 127-140.
- Dreyer, I, Antunes, S, Hoshi, T, MullerRober, B, Palme, K, Pongs, O, Reintanz, B & Hedrich, R 1997, 'Plant K<sup>+</sup> channel alpha-subunits assemble indiscriminately', *Biophysical Journal*, vol. 72, no. 5, pp. 2143-2150.
- Dreyer, I, Poree, F, Schneider, A, Mittelstadt, J, Bertl, A, Sentenac, H, Thibaud, JB & Mueller-Roeber, B 2004, 'Assembly of plant Shaker-like K-out channels requires two distinct sites of the channel alpha-subunit', *Biophysical Journal*, vol. 87, no. 2, pp. 858-872.
- Drozdowicz, YM & Rea, PA 2001, 'Vacuolar H<sup>+</sup> pyrophosphatases: from the evolutionary backwaters into the mainstream', *Trends in Plant Science*, vol. 6, no. 5, pp. 206-211.
- Dubcovsky, J, Maria, GS, Epstein, E, Luo, MC & Dvorak, J 1996, 'Mapping of the K<sup>+</sup>/Na<sup>+</sup> discrimination locus Kna1 in wheat', *Theoretical and Applied Genetics*, vol. 92, no. 3-4, pp. 448-454.
- Edwards, K, Johnstone, C & Thompson, C 1991, 'A simple and rapid method for the preparation of plant genomic DNA for PCR analysis', *Nucleic Acids Research*, vol. 19, no. 6, pp. 1349-1349.
- Epstein, E 1972, *Mineral nutrition of plants: principles and perspectives.*, London, John Wiley and Sons, Inc. UK.
- Fan, TWM, Higashi, RM, Norlyn, J & Epstein, E 1989, '*In vivo* <sup>23</sup>Na- and <sup>31</sup>P- NMR measurment of a tonoplast Na<sup>+</sup>/H<sup>+</sup> exchange process and its characteristics in two barley cultivars', *Proceedings of the National Academy of Sciences of the United States of America*, vol. 86, no. 24, pp. 9856-9860.
- FAO 2008, *Land and Plant Nutrition Managment Service. United Nation Food and Agriculture Organization*, Rome, Italy.
- Ferjani, A, Segami, S, Horiguchi, G, Muto, Y, Maeshima, M & Tsukaya, H 2011, 'Keep an eye on PPi: The vacuolar-type H<sup>+</sup>-pyrophosphatase regulates postgerminative development in Arabidopsis', *Plant Cell*, vol. 23, no. 8, pp. 2895-2908.
- Flowers, T, Galal, H & Bromham, L 2010, 'Evolution of halophytes: multiple origins of salt tolerance in land plants', *Functional Plant Biology*, vol. 37, pp. 604-612.
- Flowers, TJ 1999, 'Salinisation and horticultural production', *Scientia Horticulturae*, vol. 78, no. 1-4, pp. 1-4.

- Flowers, TJ 2004, 'Improving crop salt tolerance', *Journal of Experimental Botany*, vol. 55, no. 396, pp. 307-319.
- Flowers, TJ & Colmer, TD 2008, 'Salinity tolerance in halophytes', *New Phytologist*, vol. 179, no. 4, pp. 945-963.
- Flowers, TJ, Troke, PF & Yeo, AR 1977, 'Mechanism of salt tolerance in halophytes', *Annual Review of Plant Physiology and Plant Molecular Biology*, vol. 28, pp. 89-121.
- Flowers, TJ & Yeo, AR 1995, 'Breeding for salinity resistance in crop plants: Where next?', *Australian Journal of Plant Physiology*, vol. 22, no. 6, pp. 875-884.
- Frensch, J & Hsiao, TC 1994, 'Transient responses of cell turgor and growth of maize roots as affected by changes in water potential', *Plant Physiology*, vol. 104, no. 1, pp. 247-254.
- Frensch, J & Hsiao, TC 1995, 'Rapid response of the yield threshold and turgor regulation during adjustment of root-growth to water-stress in *Zea mays*', *Plant Physiology*, vol. 108, no. 1, pp. 303-312.
- Fukuda, A, Chiba, K, Maeda, M, Nakamura, A, Maeshima, M & Tanaka, Y 2004a, 'Effect of salt and osmotic stresses on the expression of genes for the vacuolar H<sup>+</sup>-pyrophosphatase, H<sup>+</sup>-ATPase subunit A, and Na<sup>+</sup>/H<sup>+</sup> antiporter from barley', *Journal of Experimental Botany*, vol. 55, no. 397, pp. 585-594.
- Fukuda, A, Nakamura, A, Tagiri, A, Tanaka, H, Miyao, A, Hirochika, H & Tanaka, Y 2004b, 'Function, intracellular localization and the importance in salt tolerance of a vacuolar Na<sup>+</sup>/H<sup>+</sup> antiporter from rice', *Plant and Cell Physiology*, vol. 45, no. 2, pp. 146-159.
- Fukuda, A, Yazaki, Y, Ishikawa, T, Koike, S & Tanaka, Y 1998, 'Na<sup>+</sup>/H<sup>+</sup> antiporter in tonoplast vesicles from rice roots', *Plant and Cell Physiology*, vol. 39, no. 2, pp. 196-201.
- Furuichi, T, Cunningham, KW & Muto, S 2001, 'A putative two pore channel *AtTPC1* mediates Ca<sup>2+</sup> flux in Arabidopsis leaf cells', *Plant and Cell Physiology*, vol. 42, no. 9, pp. 900-905.
- Garbarino, J & Dupont, FM 1988, 'NaCl induces a Na<sup>+</sup>/H<sup>+</sup> antiporter in tonoplast vesicles from barley roots', *Plant Physiology*, vol. 86, pp. 0231-0236.
- Garbarino, J & Dupont, FM 1989, 'Rapid induction of Na<sup>+</sup>/H<sup>+</sup> exchange activity in barley root tonoplast', *Plant Physiology*, vol. 89, no. 1, pp. 1-4.
- Garciadeblas, B, Benito, B & Rodriguez-Navarro, A 2001, 'Plant cells express several stress calcium ATPases but apparently no sodium ATPase', *Plant and Soil*, vol. 235, no. 2, pp. 181-192.
- Garciadeblas, B, Senn, ME, Banuelos, MA & Rodriguez-Navarro, A 2003, 'Sodium transport and HKT transporters: the rice model', *Plant Journal*, vol. 34, no. 6, pp. 788-801.
- Gaxiola, RA, Fink, GR & Hirschi, KD 2002, 'Genetic manipulation of vacuolar proton pumps and transporters', *Plant Physiology*, vol. 129, no. 3, pp. 967-973.

- Gaxiola, RA, Li, JS, Undurraga, S, Dang, LM, Allen, GJ, Alper, SL & Fink, GR 2001, 'Drought- and salt-tolerant plants result from overexpression of the *AVP1* H<sup>+</sup>-pump', *Proceedings of the National Academy of Sciences of the United States of America*, vol. 98, no. 20, pp. 11444-11449.
- Gaxiola, RA, Palmgren, MG & Schumacher, K 2007, 'Plant proton pumps', *FEBS Letters*, vol. 581, no. 12, pp. 2204-2214.
- Gaxiola, RA, Rao, R, Sherman, A, Grisafi, P, Alper, SL & Fink, GR 1999, 'The *Arabidopsis thaliana* proton transporters, *AtNhx1* and *Avp1*, can function in cation detoxification in yeast', *Proceedings of the National Academy of Sciences of the United States of America*, vol. 96, pp. 1480-1485.
- Gaxiola, RA, Sanchez, CA, Paez-Valencia, J, Ayre, BG & Elser, JJ 2012, 'Genetic manipulation of a "Vacuolar" H<sup>+</sup>-PPase: From salt tolerance to yield enhancement under phosphorus-deficient soils', *Plant Physiology*, vol. 159, no. 1, pp. 3-11.
- Glenn, EP, Brown, JJ & Blumwald, E 1999, 'Salt tolerance and crop potential of halophytes', *Critical Reviews in Plant Sciences*, vol. 18, no. 2, pp. 227-255.
- Gonzalez, N, De Bodt, S, Sulpice, R, Jikumaru, Y, Chae, E, Dhondt, S, Van Daele, T, De Milde, L, Weigel, D, Kamiya, Y, Stitt, M, Beemster, GTS & Inze, D 2010, 'Increased leaf size: Different means to an end', *Plant Physiology*, vol. 153, no. 3, pp. 1261-1279.
- Gorham, J, Hardy, C, Wyn Jones, RG, Joppa, LR & Law, CN 1987, 'Chromosomal location of a K<sup>+</sup>/Na<sup>+</sup> discrimination character in the D genome of wheat', *Theoretical and Applied Genetics*, vol. 74, pp. 584-588.
- Gorham, J, Jones, RGW & Bristol, A 1990, 'Partial characterization of the trait for enhanced K<sup>+</sup>/Na<sup>+</sup> discrimination in the D-genome of wheat', *Planta*, vol. 180, no. 4, pp. 590-597.
- Greenway, H & Munns, R 1980, 'Mechanism of salt tolerance in nonhalophytes', *Annual Review of Plant Physiology*, vol. 31, pp. 149-190.
- Guo, SL, Yin, HB, Zhang, X, Zhao, FY, Li, PH, Chen, SH, Zhao, YX & Zhang, H 2006, 'Molecular cloning and characterization of a vacuolar H<sup>+</sup>-pyrophosphatase gene, *SsVP*, from the halophyte *Suaeda salsa* and its overexpression increases salt and drought tolerance of *Arabidopsis*', *Plant Molecular Biology*, vol. 60, no. 1, pp. 41-50.
- Hamaji, K, Nagira, M, Yoshida, K, Ohnishi, M, Oda, Y, Uemura, T, Goh, T, Sato, MH, Morita, MT, Tasaka, M, Hasezawa, S-i, Nakano, A, Hara-Nishimura, I, Maeshima, M, Fukaki, H & Mimura, T 2009, 'Dynamic Aspects of Ion Accumulation by Vesicle Traffic Under Salt Stress in *Arabidopsis*', *Plant and Cell Physiology*, vol. 50, no. 12, pp. 2023-2033.
- Hariadi, Y, Marandon, K, Tian, Y, Jacobsen, SE & Shabala, S 2011, 'Ionic and osmotic relations in quinoa (*Chenopodium quinoa* Willd.) plants grown at various salinity levels', *Journal of Experimental Botany*, vol. 62, no. 1, pp. 185-193.
- Hasegawa, PM, Bressan, RA, Zhu, JK & Bohnert, HJ 2000, 'Plant cellular and molecular responses to high salinity', *Annual Review of Plant Physiology and Plant Molecular Biology*, vol. 51, pp. 463-499.

- Hayashi, H, Alia, Sakamoto, A, Nonaka, H, Chen, THH & Murata, N 1998, 'Enhanced germination under high-salt conditions of seeds of transgenic Arabidopsis with a bacterial gene (*codA*) for choline oxidase', *Journal of Plant Research*, vol. 111, no. 1102, pp. 357-362.
- He, XL, Huang, X, Shen, YZ & Huang, ZJ 2014, 'Wheat V-H<sup>+</sup>-ATPase subunit genes significantly affect salt tolerance in *Arabidopsis thaliana*', *Plos One*, vol. 9, no. 1.
- Hedrich, R, Barbierbrygoo, H, Felle, H, Flugge, UI, Luttge, U, Maathuis, FJM, Marx, S, Prins, HBA, Raschke, K, Schnabl, H, Schroeder, JI, Struve, I, Taiz, L & Ziegler, P 1988, 'General mechanisms for solute transport across the tonoplast of plant vacuoles-A patch clamp survey of ion channels and proton pumps', *Botanica Acta*, vol. 101, no. 1, pp. 7-13.
- Hedrich, R & Marten, I 2011, 'TPC1-SV Channels Gain Shape', *Molecular Plant*, vol. 4, no. 3, pp. 428-441.
- Hedrich, R & Neher, E 1987, 'Cytoplasmic calcium regulates voltage-dependent ion channels in plant vacuoles', *Nature*, vol. 329, no. 6142, pp. 833-836.
- Heinonen, JK 2001, *Biological role of inorganic pyrophosphate*, Kluwer Academic Publishers, London, United Kingdom.
- Holmstrom, KO, Somersalo, S, Mandal, A, Palva, TE & Welin, B 2000, 'Improved tolerance to salinity and low temperature in transgenic tobacco producing glycine betaine', *Journal of Experimental Botany*, vol. 51, no. 343, pp. 177-185.
- Hughes, FM & Cidlowski, JA 1998, 'Glucocorticoid-induced thymocyte apoptosis: Protease-dependent activation of cell shrinkage and DNA degradation', *Journal of Steroid Biochemistry and Molecular Biology*, vol. 65, no. 1-6, pp. 207-217.
- Hughes, FM & Cidlowski, JA 1999, 'Potassium is a critical regulator of apoptotic enzymes in vitro and in vivo', in G Weber (ed.), *Advances in Enzyme Regulation, Vol 39*, vol. 39, pp. 157-171.
- Inan, G, Zhang, Q, Li, PH, Wang, ZL, Cao, ZY, Zhang, H, Zhang, CQ, Quist, TM, Goodwin, SM, Zhu, JH, Shi, HH, Damsz, B, Charbaji, T, Gong, QQ, Ma, SS, Fredricksen, M, Galbraith, DW, Jenks, MA, Rhodes, D, Hasegawa, PM, Bohnert, HJ, Joly, RJ, Bressan, RA & Zhu, JK 2004, 'Salt cress. A halophyte and cryophyte Arabidopsis relative model system and its applicability to molecular genetic analyses of growth and development of extremophiles', *Plant Physiology*, vol. 135, no. 3, pp. 1718-1737.
- Isayenkov, S, Isner, JC & Maathuis, FJM 2010, 'Vacuolar ion channels: Roles in plant nutrition and signalling', *FEBS Letters*, vol. 584, no. 10, pp. 1982-1988.
- Ivashikina, N & Hedrich, R 2005, 'K<sup>+</sup> currents through SV-type vacuolar channels are sensitive to elevated luminal sodium levels', *Plant Journal*, vol. 41, no. 4, pp. 606-614.
- Jacobs, A, Lunde, C, Bacic, A, Tester, M & Roessner, U 2007, 'The impact of constitutive heterologous expression of a moss Na<sup>+</sup> transporter on the metabolomes of rice and barley', *Metabolomics*, vol. 3, no. 3, pp. 307-317.



- Jacobsen, SE, Mujica, A & Jensen, CR 2003, 'The resistance of quinoa (*Chenopodium quinoa* Willd.) to adverse abiotic factors', *Food Reviews International*, vol. 19, no. 1-2, pp. 99-109.
- Jeschke, WD, Stelter, W, Reising, B & Behl, R 1983, 'Vacuolar  $\text{Na}^+/\text{K}^+$  exchange, its occurrence in root-cells of *Hordeum*, *Atriplex* and *Zea* and its significance for  $\text{K}^+/\text{Na}^+$  discrimination in roots', *Journal of Experimental Botany*, vol. 34, no. 145, pp. 964-979.
- Kawano, T 2003, 'Roles of the reactive oxygen species-generating peroxidase reactions in plant defense and growth induction', *Plant Cell Reports*, vol. 21, no. 9, pp. 829-837.
- Keller, A, Nesvizhskii, AI, Kolker, E & Aebersold, R 2002, 'Empirical statistical model to estimate the accuracy of peptide identifications made by MS/MS and database search', *Analytical Chemistry*, vol. 74, no. 20, pp. 5383-5392.
- Krebs, M, Beyhl, D, Gorlich, E, Al-Rasheid, KAS, Marten, I, Stierhof, YD, Hedrich, R & Schumacher, K 2010, 'Arabidopsis V-ATPase activity at the tonoplast is required for efficient nutrient storage but not for sodium accumulation', *Proceedings of the National Academy of Sciences of the United States of America*, vol. 107, no. 7, pp. 3251-3256.
- Kurusu, T, Sakurai, Y, Miyao, A, Hirochika, H & Kuchitsu, K 2004, 'Identification of a putative voltage-gated  $\text{Ca}^{2+}$ -permeable channel (*OsTPC1*) involved in  $\text{Ca}^{2+}$  influx and regulation of growth and development in rice', *Plant and Cell Physiology*, vol. 45, no. 6, pp. 693-702.
- Leidi, OE, Barragan, V, Rubio, L, El-Hamdaoui, A, Ruiz, TM, Cubero, B, Fernandez, AJ, Bressan, AR, Hasagawa, MP, Quintero, JF & Pardo, MJ 2010, 'The *AtNHX1* exchanger mediates potassium compartmentation in vacuoles of transgenic tomato', *The Plant Journal*, vol. 61, pp. 495-506.
- Li, JS, Yang, HB, Peer, WA, Richter, G, Blakeslee, J, Bandyopadhyay, A, Titapiwantakun, B, Undurraga, S, Khodakovskaya, M, Richards, EL, Krizek, B, Murphy, AS, Gilroy, S & Gaxiola, R 2005, 'Arabidopsis  $\text{H}^+$ -PPase *AVP1* regulates auxin-mediated organ development', *Science*, vol. 310, no. 5745, pp. 121-125.
- Lilius, G, Holmberg, N & Bulow, L 1996, 'Enhanced NaCl stress tolerance in transgenic tobacco expressing bacterial choline dehydrogenase', *Bio-Technology*, vol. 14, no. 2, pp. 177-180.
- Lu, Z, Percy, R, Qualset, C & Zeiger, E 1998, 'Stomatal conductance predicts yields in irrigated Pima cotton and bread wheat grown at high temperatures', *Journal of Experimental Botany*, vol. 49, no. Special Issue, pp. 453-460.
- Maathuis, FJM 2014, 'Sodium in plants: perception, signalling, and regulation of sodium fluxes', *Journal of Experimental Botany*, vol. 65, no. 3, pp. 849-858.
- Maathuis, FJM & Amtmann, A 1999, ' $\text{K}^+$  nutrition and  $\text{Na}^+$  toxicity: The basis of cellular  $\text{K}^+/\text{Na}^+$  ratios', *Annals of Botany*, vol. 84, pp. 123-133.
- Maathuis, FJM, Flowers, TJ & Yeo, AR 1992, 'Sodium-chloride compartmentation in leaf vacuoles of the halophyte *Suaeda maritima* (L) Dum and its relation to tonoplast permeability', *Journal of Experimental Botany*, vol. 43, no. 254, pp. 1219-1223.

- Maathuis, FJM & Prins, HBA 1990, 'Patch clamp studies on root cell vacuoles of salt tolerant and salt sensitive *Plantago* species- regulation of channel activity by salt stress', *Plant Physiology*, vol. 92, no. 1, pp. 23-28.
- Maeshima, M 2000, 'Vacuolar H<sup>+</sup>-pyrophosphatase', *Biochimica et Biophysica Acta (BBA) - Biomembranes*, vol. 1465 no. 1-2, pp. 37-51.
- Mahajan, S & Tuteja, N 2005, 'Cold, salinity and drought stresses: An overview', *Archives of Biochemistry and Biophysics*, vol. 444, no. 2, pp. 139-158.
- Maksimovic, JD, Zhang, JY, Zeng, FR, Zivanovic, BD, Shabala, L, Zhou, MX & Shabala, S 2013, 'Linking oxidative and salinity stress tolerance in barley: can root antioxidant enzyme activity be used as a measure of stress tolerance?', *Plant and Soil*, vol. 365, no. 1-2, pp. 141-155.
- Matoh, T, Ishikawa, T & Takahashi, E 1989, 'Collapse of ATP-induced pH gradient by sodium ions in microsomal membrane vesicles prepared from *Atriplex gemlini* leaves- possibility of Na<sup>+</sup>/H<sup>+</sup> antiport', *Plant Physiology*, vol. 89, no. 1, pp. 180-183.
- Mengel, K & Arneke, W-W 1982, 'Effect of potassium on water potential, the pressure potential, the osmotic potential and cell elongation in leaves of *Phaseolus vulgaris*', *Physiologia Plantarum*, vol. 54, pp. 402-408.
- Mian, A 2010, 'Improving salt stress resistance in cereals', PhD thesis, University of York.
- Mian, A, Oomen, R, Isayenkov, S, Sentenac, H, Maathuis, FJM & Very, AA 2011, 'Over-expression of an Na<sup>+</sup> and K<sup>+</sup> permeable *HKT* transporter in barley improves salt tolerance', *Plant Journal*, vol. 68, no. 3, pp. 468-479.
- Millar, J & Roots, J 2012, 'Changes in Australian agriculture and land use: implications for future food security', *International Journal of Agricultural Sustainability*, vol. 10, no. 1, pp. 25-39.
- Miller, G, Shulaev, V & Mittler, R 2008, 'Reactive oxygen signaling and abiotic stress', *Physiologia Plantarum*, vol. 133, no. 3, pp. 481-489.
- Mittler, R 2002, 'Oxidative stress, antioxidants and stress tolerance', *Trends in Plant Science*, vol. 7, no. 9, pp. 405-410.
- Moller, IS, Gilliam, M, Jha, D, Mayo, GM, Roy, SJ, Coates, JC, Haseloff, J & Tester, M 2009, 'Shoot Na<sup>+</sup> exclusion and increased salinity tolerance engineered by cell type-specific alteration of Na<sup>+</sup> transport in *Arabidopsis*', *Plant Cell*, vol. 21, no. 7, pp. 2163-2178.
- Munns, R 1993, 'Physiological processes limiting plant growth in saline soils-some dogmas and hypotheses', *Plant Cell and Environment*, vol. 16, no. 1, pp. 15-24.
- Munns, R 2002, 'Comparative physiology of salt and water stress', *Plant, Cell & Environment*, vol. 25, pp. 239-250.
- Munns, R & Gilliam, M 2015, 'Salinity tolerance of crops - what is the cost?', *New Phytologist*.

- Munns, R, James, RA, Xu, B, Athman, A, Conn, SJ, Jordans, C, Byrt, CS, Hare, RA, Tyerman, SD, Tester, M, Plett, D & Gilliam, M 2012, 'Wheat grain yield on saline soils is improved by an ancestral Na<sup>+</sup> transporter gene', *Nature Biotechnology*, vol. 30, no. 4, pp. 360-U173.
- Munns, R, Schachtman, DP & Condon, AG 1995, 'The significance of a two-phase growth response to salinity in wheat and barley', *Australian Journal of Plant Physiology*, vol. 22, no. 4, pp. 561-569.
- Munns, R & Termaat, A 1986, 'Whole-plant responses to salinity', *Australian Journal of Plant Physiology*, vol. 13, no. 1, pp. 143-160.
- Munns, R & Tester, M 2008, 'Mechanisms of salinity tolerance', *Annual Review of Plant Biology*, vol. 59, pp. 651-681.
- Nakanishi, Y & Maeshima, M 1998, 'Molecular cloning of vacuolar H<sup>+</sup>-pyrophosphatase and its developmental expression in growing hypocotyl of mung bean', *Plant Physiology*, vol. 116, no. 2, pp. 589-597.
- Nass, R, Cunningham, KW & Rao, R 1997, 'Intracellular sequestration of sodium by a novel Na<sup>+</sup>/H<sup>+</sup> exchanger in yeast is enhanced by mutations in the plasma membrane H<sup>+</sup>-ATPase - Insights into mechanisms of sodium tolerance', *Journal of Biological Chemistry*, vol. 272, no. 42, pp. 26145-26152.
- Nass, R & Rao, R 1998, 'Novel localization of a Na<sup>+</sup>/H<sup>+</sup> exchanger in a late endosomal compartment of yeast - Implications for vacuole biogenesis', *Journal of Biological Chemistry*, vol. 273, no. 33, pp. 21054-21060.
- Nesvizhskii, AI, Keller, A, Kolker, E & Aebersold, R 2003, 'A Statistical Model for Identifying Proteins by Tandem Mass Spectrometry', *Analytical Chemistry*, vol. 75, no. 17, pp. 4646-4658.
- Neuhaus, HE & Trentmann, O 2014, 'Regulation of transport processes across the tonoplast', *Frontiers in Plant Science*, vol. 5.
- Nieves-Cordones, M, Miller, AJ, Aleman, F, Martinez, V & Rubio, F 2008, 'A putative role for the plasma membrane potential in the control of the expression of the gene encoding the tomato high-affinity potassium transporter *HAK5*', *Plant Molecular Biology*, vol. 68, no. 6, pp. 521-532.
- Ohta, M, Hayashi, Y, Nakashima, A, Hamada, A, Tanaka, A, Nakamura, T & Hayakawa, T 2002, 'Introduction of a Na<sup>+</sup>/H<sup>+</sup> antiporter gene from *Atriplex gmelini* confers salt tolerance to rice', *FEBS Letters*, vol. 532, no. 3, pp. 279-282.
- Padmanaban, S, Lin, XY, Perera, I, Kawamura, Y & Sze, H 2004, 'Differential expression of vacuolar H<sup>+</sup>-ATPase subunit c genes in tissues active in membrane trafficking and their roles in plant growth as revealed by RNAi', *Plant Physiology*, vol. 134, no. 4, pp. 1514-1526.
- Palmgren, MG & Nissen, P 2011, 'P-Type ATPases', in DC Rees, KA Dill & JR Williamson (eds), *Annual Review of Biophysics*, Vol 40, vol. 40, pp. 243-266.
- Panta, S, Flowers, T, Lane, P, Doyle, R, Haros, G & Shabala, S 2014, 'Halophyte agriculture: Success stories', *Environmental and Experimental Botany*, vol. 107, pp. 71-83.

- Pantoja, O, Dainty, J & Blumwald, E 1989, 'Ion channels in vacuoles from halophytes and glycophytes', *FEBS Letters*, vol. 255, no. 1, pp. 92-96.
- Pardo, JM, Cubero, B, Leidi, EO & Quintero, FJ 2006, 'Alkali cation exchangers: roles in cellular homeostasis and stress tolerance', *Journal of Experimental Botany*, vol. 57, no. 5, pp. 1181-1199.
- Parks, GE, Dietrich, MA & Schumaker, KS 2002, 'Increased vacuolar  $\text{Na}^+/\text{H}^+$  exchange activity in *Salicornia bigelovii* Torr. in response to NaCl', *Journal of Experimental Botany*, vol. 53, no. 371, pp. 1055-1065.
- Passioura, JB & Munns, R 2000, 'Rapid environmental changes that affect leaf water status induce transient surges or pauses in leaf expansion rate', *Australian Journal of Plant Physiology*, vol. 27, no. 10, pp. 941-948.
- Peiter, E, Maathuis, FJM, Mills, LN, Knight, H, Hetherington, AM, Pelloux, J & Sanders, D 2005, 'TPC1, a vacuolar  $\text{Ca}^{2+}$ -activated channel controlling germination and stomatal movement', *Comparative Biochemistry and Physiology a-Molecular & Integrative Physiology*, vol. 141, no. 3, pp. S345-S345.
- Perez-Castineira, JR, Hernandez, A, Drake, R & Serrano, A 2011, 'A plant proton-pumping inorganic pyrophosphatase functionally complements the vacuolar ATPase transport activity and confers bafilomycin resistance in yeast', *Biochemical Journal*, vol. 437, pp. 269-278.
- Perez-Sayans, M, Suarez-Penaranda, JM, Barros-Angueira, F, Diz, PG, Gandara-Rey, JM & Garcia-Garcia, A 2012, 'An update in the structure, function, and regulation of V-ATPases: the role of the C subunit', *Brazilian Journal of Biology*, vol. 72, no. 1, pp. 189-198.
- Pittman, JK 2012, 'Multiple transport pathways for mediating intracellular pH homeostasis: the contribution of  $\text{H}^+$ /ion exchangers', *Frontiers in Plant Science*, vol. 3, no. JAN, pp. 1-8.
- Plett, D, Safwat, G, Gilliam, M, Møller, I, Roy, S, Shirley, N, Jacobs, A, Johnson, A & Tester, M 2010a, 'Improved salinity tolerance of rice through cell type-specific expression of *AtHKT1;1*', *Plos One*, vol. 5, no. 9, pp. 1-8.
- Plett, DC & Moller, IS 2010b, ' $\text{Na}^+$  transport in glycophytic plants: what we know and would like to know', *Plant Cell and Environment*, vol. 33, no. 4, pp. 612-626.
- Pottosin, II, Dobrovinskaya, OR & Muniz, J 2001, 'Conduction of monovalent and divalent cations in the slow vacuolar channel', *Journal of Membrane Biology*, vol. 181, no. 1, pp. 55-65.
- Pottosin, II, Tikhonova, LI, Hedrich, R & Schonknecht, G 1997, 'Slowly activating vacuolar channels can not mediate  $\text{Ca}^{2+}$ -induced  $\text{Ca}^{2+}$  release', *Plant Journal*, vol. 12, no. 6, pp. 1387-1398.
- Pottosin, I & Dobrovinskaya, O 2014, 'Non-selective cation channels in plasma and vacuolar membranes and their contribution to  $\text{K}^+$  transport', *Journal of Plant Physiology*, vol. 171 no. 9, pp. 732-742.
- Pottosin, I, Martinez-Estevéz, M, Dobrovinskaya, O, Muniz, J & Schonknecht, G 2004, 'Mechanism of luminal  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  action on the vacuolar slowly activating channels', *Planta*, vol. 219, no. 6, pp. 1057-1070.

- Poustini, K & Siosemardeh, A 2004, 'Ion distribution in wheat cultivars in response to salinity stress', *Field Crops Research*, vol. 85, no. 2-3, pp. 125-133.
- Qadir, M, Quill  rou, E, Nangia, V, Murtaza, G, Singh, M, Thomas, RJ, Drechsel, P & Noble, AD 2014, 'Economics of salt-induced land degradation and restoration', *Natural Resources Forum*, vol. 38, no. 4, pp. 282-295.
- Qadir, M, Tubeileh, A, Akhtar, J, Larbi, A, Minhas, P & Khan, M 2008, 'Productivity enhancement of salt-affected environments through crop diversification', *Land Degradation and Development*, vol. 19, pp. 429-453.
- Qiu, NW, Chen, M, Guo, JR, Bao, HY, Ma, XL & Wang, BS 2007, 'Coordinate up-regulation of V-H<sup>+</sup>-ATPase and vacuolar Na<sup>+</sup>/H<sup>+</sup> antiporter as a response to NaCl treatment in a C-3 halophyte *Suaeda salsa*', *Plant Science*, vol. 172, no. 6, pp. 1218-1225.
- Qiu, Q-S 2012, 'Plant and yeast NHX antiporters: Roles in membrane trafficking', *Journal of Integrative Plant Biology*, vol. 54, no. 2, pp. 66-72.
- Qiu, Q-S, Guo, Y, Dietrich, AM, Schumaker, SK & Zhu, J-K 2002, 'Regulation of SOS1, a plasma membrane Na<sup>+</sup>/H<sup>+</sup> exchanger in *Arabidopsis thaliana*, by SOS2 and SOS3', *Proceedings of the National Academy of Sciences of the United States of America*, vol. 99, no. 12, pp. 8436-8844.
- Qiu, Q-S, Guo, Y, Quintero, JF, Pardo, MJ & Schumaker, SK 2004, 'Regulation of Vacuolar Na/H Exchange in *Arabidopsis thaliana* by the Salt-Overly-Sensitive (SOS) Pathway', *Journal of Biological Chemistry*, vol. 279, no. 1, pp. 207-215.
- Qiu, QS, Barkla, BJ, Vera-Estrella, R, Zhu, JK & Schumaker, KS 2003, 'Na<sup>+</sup>/H<sup>+</sup> exchange activity in the plasma membrane of *Arabidopsis*', *Plant Physiology*, vol. 132, no. 2, pp. 1041-1052.
- Queiros, F, Fontes, N, Silva, P, Almeida, D, Maeshima, M, Geros, H & Fidalgo, F 2009, 'Activity of tonoplast proton pumps and Na<sup>+</sup>/H<sup>+</sup> exchange in potato cell cultures is modulated by salt', *Journal of Experimental Botany*, vol. 60, no. 4, pp. 1363-1374.
- Quintero, FJ, Blatt, MR & Pardo, JM 2000, 'Functional conservation between yeast and plant endosomal Na<sup>+</sup>/H<sup>+</sup> antiporters', *FEBS Letters*, vol. 471, no. 2-3, pp. 224-228.
- Quintero, FJ, Martinez-Atienza, J, Villalta, I, Jiang, XY, Kim, WY, Ali, Z, Fujii, H, Mendoza, I, Yun, DJ, Zhu, JK & Pardo, JM 2011, 'Activation of the plasma membrane Na/H antiporter Salt-Overly-Sensitive 1 (SOS1) by phosphorylation of an auto-inhibitory C-terminal domain', *Proceedings of the National Academy of Sciences of the United States of America*, vol. 108, no. 6, pp. 2611-2616.
- Rajendran, K, Tester, M & Roy, SJ 2009, 'Quantifying the three main components of salinity tolerance in cereals', *Plant Cell and Environment*, vol. 32, no. 3, pp. 237-249.
- Rauch, A, Bellew, M, Eng, J, Fitzgibbon, M, Holzman, T, Hussey, P, Igra, M, MacLean, B, Lin, CW, Detter, A, Fang, R, Faca, V, Gafken, P, Zhang, H, Whiteaker, J, States, D, Hanash, S, Paulovich, A & McIntosh, MW 2006, 'Computational Proteomics Analysis System (CPAS): An

- Extensible, Open-Source Analytic System for Evaluating and Publishing Proteomic Data and High Throughput Biological Experiments', *Journal of Proteome Research*, vol. 5, no. 4, pp. 1024-1024.
- Raven, JA 1985, 'Regulation of pH and generation of osmolarity in vascular plants-A cost-benefit analysis in relation to efficiency of use of energy, nitrogen and water', *New Phytologist*, vol. 101, no. 1, pp. 25-77.
- Rea, PA & Poole, RJ 1993, 'Vacuolar H<sup>+</sup>-translocating pyrophosphatase', *Annual Review of Plant Physiology and Plant Molecular Biology*, vol. 44, pp. 157-180.
- Ren, ZH, Gao, JP, Li, LG, Cai, XL, Huang, W, Chao, DY, Zhu, MZ, Wang, ZY, Luan, S & Lin, HX 2005, 'A rice quantitative trait locus for salt tolerance encodes a sodium transporter', *Nature Genetics*, vol. 37, no. 10, pp. 1141-1146.
- Rengasamy, P 2002, 'Transient salinity and subsoil constraints to dryland farming in Australian sodic soils: an overview', *Australian Journal of Experimental Agriculture*, vol. 42, pp. 351-361.
- Rengasamy, P 2006, 'World salinization with emphasis on Australia', *Journal of Experimental Botany*, vol. 57, pp. 1017-1023.
- Riadh, K, Wided, M, Hans-Werner, K & Chedly, A 2010, 'Responses of halophytes to environmental stresses with special emphasis to salinity', in JC Kader & M Delseny (eds), *Advances in Botanical Research, Vol 53*, vol. 53, pp. 117-145.
- Richards, RA 2000, 'Selectable traits to increase crop photosynthesis and yield of grain crops', *Journal of Experimental Botany*, vol. 51, no. GMP Special Issue, pp. 447-458.
- Rodriguez-Rosales, MP, Jiang, XY, Galvez, FJ, Aranda, MN, Cubero, B & Venema, K 2008, 'Overexpression of the tomato K<sup>+</sup>/H<sup>+</sup> antiporter *LeNHX2* confers salt tolerance by improving potassium compartmentalization', *New Phytologist*, vol. 179, no. 2, pp. 366-377.
- Rodriguez, HG, Roberts, JKM, Jordan, WR & Drew, MC 1997, 'Growth, water relations, and accumulation of organic and inorganic solutes in roots of maize seedlings during salt stress', *Plant Physiology*, vol. 113, no. 3, pp. 881-893.
- Roy, SJ, Huang, W, Wang, XJ, Evrard, A, Schmockel, SM, Zafar, ZU & Tester, M 2013, 'A novel protein kinase involved in Na<sup>+</sup> exclusion revealed from positional cloning', *Plant Cell and Environment*, vol. 36, no. 3, pp. 553-568.
- Ruan, C, da Silva, J, Mopper, S, Qin, P & Lutts, S 2010, 'Halophyte improvment for a salinized world', *Critical Reviews in Plant Sciences*, vol. 29, pp. 329-359.
- Sagi, M & Fluhr, R 2006, 'Production of reactive oxygen species by plant NADPH oxidases', *Plant Physiology*, vol. 141, no. 2, pp. 336-340.
- Sander, JD & Joung, JK 2014, 'CRISPR-Cas systems for editing, regulating and targeting genomes', *Nature Biotechnology*, vol. 32, no. 4, pp. 347-355.

- Schachtman, DP, Bloom, AJ & Dvorak, J 1989, 'Salt-tolerant Triticum x Lophopyrum derivatives limit the accumulation of sodium and chloride ions under saline-stress', *Plant, Cell & Environment*, vol. 12, no. 1, pp. 47-55.
- Schilling, RK, Marschner, P, Shavrukov, Y, Berger, B, Tester, M, Roy, SJ & Plett, DC 2014, 'Expression of the Arabidopsis vacuolar H<sup>+</sup> - pyrophosphatase gene (*AVPI*) improves the shoot biomass of transgenic barley and increases grain yield in a saline field', *Plant Biotechnology Journal*, vol. 12, no. 3, pp. 378-386.
- Scholz-Starke, J, De Angeli, A, Ferraretto, C, Paluzzi, S, Gambale, F & Carpaneto, A 2004, 'Redox-dependent modulation of the carrot SV channel by cytosolic pH', *FEBS Letters*, vol. 576, no. 3, pp. 449-454.
- Schulzlessdorf, B & Hedrich, R 1995, 'Protons and calcium modulate SV-type channels in the vacuolar-lysosomal compartment- channel interaction with calmodulin inhibitors', *Planta*, vol. 197, no. 4, pp. 655-671.
- Schumacher, K & Krebs, M 2010, 'The V-ATPase: small cargo, large effects', *Current Opinion in Plant Biology*, vol. 13, no. 6, pp. 724-730.
- Segami, S, Nakanishi, Y, Sato, MH & Maeshima, M 2010, 'Quantification, organ-specific accumulation and intracellular localization of Type II H<sup>+</sup>-Pyrophosphatase in Arabidopsis thaliana', *Plant and Cell Physiology*, vol. 51, no. 8, pp. 1350-1360.
- Serraj, R & Sinclair, TR 2002, 'Osmolyte accumulation: can it really help increase crop yield under drought conditions?', *Plant Cell and Environment*, vol. 25, no. 2, pp. 333-341.
- Serrano, R & Rodriguez-Navarro, A 2001, 'Ion homeostasis during salt stress in plants', *Current Opinion in Cell Biology*, vol. 13, no. 4, pp. 399-404.
- Shabala, S 2000, 'Tonic and osmotic components of salt stress specifically modulate net ion fluxes from bean leaf mesophyll', *Plant Cell and Environment*, vol. 23, no. 8, pp. 825-837.
- Shabala, S 2009, 'Salinity and programmed cell death: unravelling mechanisms for ion specific signalling', *Journal of Experimental Botany*, vol. 60, no. 3, pp. 709-711.
- Shabala, S 2013, 'Learning from halophytes: physiological basis and strategies to improve abiotic stress tolerance in crops', *Annals of Botany*, vol. 112, no. 7, pp. 1209-1221.
- Shabala, S & Cuin, TA 2008, 'Potassium transport and plant salt tolerance', *Physiologia Plantarum*, vol. 133, no. 4, pp. 651-669.
- Shabala, S, Cuin, TA, Prismall, L & Nemchinov, LG 2007, 'Expression of animal CED-9 anti-apoptotic gene in tobacco modifies plasma membrane ion fluxes in response to salinity and oxidative stress', *Planta*, vol. 227, no. 1, pp. 189-197.
- Shabala, S, Demidchik, V, Shabala, L, Cuin, TA, Smith, SJ, Miller, AJ, Davies, JM & Newman, IA 2006, 'Extracellular Ca<sup>2+</sup> ameliorates NaCl-induced K<sup>+</sup> loss from Arabidopsis root and leaf cells by controlling plasma membrane K<sup>+</sup> permeable channels', *Plant Physiology*, vol. 141, no. 4, pp. 1653-1665.

- Shabala, S & Mackay, A 2011a, 'Ion transport in halophytes', in I.Turkan (ed.), *Plant responses to drought and salinity stress: Developments in a Post-Genomic Era*, Elsevier Ltd., vol. 57, pp. 151-199.
- Shabala, S & Munns, R 2012, *Salinity stress: physiological constraints and adaptive mechanisms*, Plant stress Physiology, CAB, UK.
- Shabala, S & Pottosin, I 2014, 'Regulation of potassium transport in plants under hostile conditions: implications for abiotic and biotic stress tolerance', *Physiologia Plantarum*, vol. 151, no. 3, pp. 257-279.
- Shabala, S & Shabala, L 2011b, 'Ion transport and osmotic adjustment in plants and bacteria', *Biomolecular Concept*, vol. 2, pp. 407-419.
- Shabala, S, Shabala, L & Van Volkenburgh, E 2003, 'Effect of calcium on root development and root ion fluxes in salinised barley seedlings', *Functional Plant Biology*, vol. 30, no. 5, pp. 507-514.
- Shabala, S, Shabala, S, Cuin, TA, Pang, JY, Percey, W, Chen, ZH, Conn, S, Eing, C & Wegner, LH 2010, 'Xylem ionic relations and salinity tolerance in barley', *Plant Journal*, vol. 61, no. 5, pp. 839-853.
- Shan, QW, Wang, YP, Li, J, Zhang, Y, Chen, KL, Liang, Z, Zhang, K, Liu, JX, Xi, JJ, Qiu, JL & Gao, CX 2013, 'Targeted genome modification of crop plants using a CRISPR-Cas system', *Nature Biotechnology*, vol. 31, no. 8, pp. 686-688.
- Shannon, MC 1997, 'Adaptation of plants to salinity', in DL Sparks (ed.), *Advances in Agronomy*, Vol 60, vol. 60, pp. 75-120.
- Shavrukov, Y, Bovill, J, Afzal, I, Hayes, JE, Roy, SJ, Tester, M & Collins, NC 2013, 'HVP10 encoding V-PPase is a prime candidate for the barley *HvNax3* sodium exclusion gene: evidence from fine mapping and expression analysis', *Planta*, vol. 237, no. 4, pp. 1111-1122.
- Shi, HZ, Ishitani, M, Kim, CS & Zhu, JK 2000, 'The *Arabidopsis thaliana* salt tolerance gene *SOS1* encodes a putative  $\text{Na}^+/\text{H}^+$  antiporter', *Proceedings of the National Academy of Sciences of the United States of America*, vol. 97, no. 12, pp. 6896-6901.
- Shi, HZ, Lee, BH, Wu, SJ & Zhu, JK 2003, 'Overexpression of a plasma membrane  $\text{Na}^+/\text{H}^+$  antiporter gene improves salt tolerance in *Arabidopsis thaliana*', *Nature Biotechnology*, vol. 21, no. 1, pp. 81-85.
- Shi, HZ, Quintero, FJ, Pardo, JM & Zhu, JK 2002a, 'The putative plasma membrane  $\text{Na}^+/\text{H}^+$  antiporter *SOS1* controls long-distance  $\text{Na}^+$  transport in plants', *Plant Cell*, vol. 14, no. 2, pp. 465-477.
- Shi, HZ & Zhu, JK 2002b, 'Regulation of expression of the vacuolar  $\text{Na}^+/\text{H}^+$  antiporter gene *AtNHX1* by salt stress and abscisic acid', *Plant Molecular Biology*, vol. 50, no. 3, pp. 543-550.



- Silva, P, Facanha, AR, Tavares, RM & Geros, H 2010, 'Role of tonoplast proton pumps and  $\text{Na}^+/\text{H}^+$  antiport system in salt tolerance of *Populus euphratica* Oliv.', *Journal of Plant Growth Regulation*, vol. 29, no. 1, pp. 23-34.
- Singh, RR, Kemp, JA, Kollmorgen, JF, Qureshi, JA & Fincher, GB 1997, 'Fertile plant regeneration from cell suspension and protoplast cultures of barley (*Hordeum vulgare* cv. Schooner)', *Plant Cell Tissue and Organ Culture*, vol. 49, no. 2, pp. 121-127.
- Smethurst, CF, Rix, K, Garnett, T, Auricht, G, Bayart, A, Lane, P, Wilson, SJ & Shabala, S 2008, 'Multiple traits associated with salt tolerance in lucerne: revealing the underlying cellular mechanisms', *Functional Plant Biology*, vol. 35, no. 7, pp. 640-650.
- Szabolcs, I 1994, 'Soils and salinisation', in M Pessarakali (ed.), *Handbook of Plant and Crop Stress*, Marcel Dekker, New York., pp. 3-11.
- Sze, H, Li, XH & Palmgren, MG 1999, 'Energization of plant cell membranes by  $\text{H}^+$ -pumping ATPases: Regulation and biosynthesis', *Plant Cell*, vol. 11, no. 4, pp. 677-689.
- Sze, H, Schumacher, K, Muller, M, Padmanaban, S & Taiz, L 2002, 'A simple nomenclature for a complex proton pump: *VHA* genes encode the vacuolar  $\text{H}^+$ -ATPase', *Trends in Plant Science*, vol. 7, no. 4, pp. 157-161.
- Termaat, A, Passioura, JB & Munns, R 1985, 'Shoot turgor does not limit shoot growth of NaCl affected wheat and barley', *Plant Physiology*, vol. 77, no. 4, pp. 869-872.
- Tester, M & Devenport, R 2003, 'Mechanism of salinity tolerance:  $\text{Na}^+$  tolerance and  $\text{Na}^+$  transport in higher plants', *Annals of Botany*, vol. 91, pp. 503-527.
- Tikhonova, LI, Pottosin, II, Dietz, KJ & Schonknecht, G 1997, 'Fast-activating cation channel in barley mesophyll vacuoles: Inhibition by calcium', *Plant Journal*, vol. 11, no. 5, pp. 1059-1070.
- Tingay, S, McElroy, D, Kalla, R, Fieg, S, Wang, MB, Thornton, S & Brettell, R 1997, 'Agrobacterium tumefaciens-mediated barley transformation', *Plant Journal*, vol. 11, no. 6, pp. 1369-1376.
- Tomos, AD, Leigh, RA, Shaw, CA & Jones, RGW 1984, 'A comparison of methods for measuring turgor pressures and osmotic pressures of cells of red beet storage tissue', *Journal of Experimental Botany*, vol. 35, no. 160, pp. 1675-1683.
- Valentine, J, Clifton-Brown, J, Hastings, A, Robson, P, Allison, G & Smith, P 2012, 'Food vs. fuel: the use of land for lignocellulosic 'next generation' energy crops that minimize competition with primary food production', *Global change Biology Bioenergy*, vol. 4, pp. 1-19.
- van den Wijngaard, PWJ, Bunney, TD, Roobeek, I, Schonknecht, G & de Boer, AH 2001, 'Slow vacuolar channels from barley mesophyll cells are regulated by 14-3-3 proteins', *FEBS Letters*, vol. 488, no. 1-2, pp. 100-104.
- Venema, K, Quintero, FJ, Pardo, JM & Donaire, JP 2002, 'The Arabidopsis  $\text{Na}^+/\text{H}^+$  exchanger *AtNHX1* catalyzes low affinity  $\text{Na}^+$  and  $\text{K}^+$  transport in reconstituted liposomes', *Journal of Biological Chemistry*, vol. 277, no. 4, pp. 2413-2418.

- Vera-Estrella, R, Barkla, BJ, Bohnert, HJ & Pantoja, O 1999, 'Salt stress in *Mesembryanthemum crystallinum* L cell suspensions activates adaptive mechanisms similar to those observed in the whole plant', *Planta*, vol. 207, no. 3, pp. 426-435.
- Vera-Estrella, R, Barkla, BJ, Garcia-Ramirez, L & Pantoja, O 2005, 'Salt stress in *Thellungiella halophila* activates  $\text{Na}^+$  transport mechanisms required for salinity tolerance', *Plant Physiology*, vol. 139, no. 3, pp. 1507-1517.
- Walker, DJ, Leigh, RA & Miller, AJ 1996, 'Potassium homeostasis in vacuolate plant cells', *Proceedings of the National Academy of Sciences of the United States of America*, vol. 93, no. 19, pp. 10510-10514.
- Wang, B, Davenport, RJ, Volkov, V & Amtmann, A 2006, 'Low unidirectional sodium influx into root cells restricts net sodium accumulation in *Thellungiella halophila*, a salt-tolerant relative of *Arabidopsis thaliana*', *Journal of Experimental Botany*, vol. 57, no. 5, pp. 1161-1170.
- Wang, BS, Luttge, U & Ratajczak, R 2001, 'Effects of salt treatment and osmotic stress on V-ATPase and V-PPase in leaves of the halophyte *Suaeda salsa*', *Journal of Experimental Botany*, vol. 52, no. 365, pp. 2355-2365.
- Wang, CM, Zhang, JL, Liu, XS, Li, Z, Wu, GQ, Cai, JY, Flowers, TJ & Wang, SM 2009, '*Puccinellia tenuiflora* maintains a low  $\text{Na}^+$  level under salinity by limiting unidirectional  $\text{Na}^+$  influx resulting in a high selectivity for  $\text{K}^+$  over  $\text{Na}^+$ ', *Plant Cell and Environment*, vol. 32, no. 5, pp. 486-496.
- Wang, L, He, XL, Zhao, YJ, Shen, YZ & Huang, ZJ 2011, 'Wheat vacuolar  $\text{H}^+$ -ATPase subunit B cloning and its involvement in salt tolerance', *Planta*, vol. 234, no. 1, pp. 1-7.
- Wang, WX, Vinocur, B & Altman, A 2003, 'Plant responses to drought, salinity and extreme temperatures: towards genetic engineering for stress tolerance', *Planta*, vol. 218, no. 1, pp. 1-14.
- Ward, JM & Schroeder, JI 1994, 'Calcium-activated  $\text{K}^+$  channels and calcium-induced calcium release by slow vacuolar ion channels in guard cell vacuoles implicated in the control of stomatal closure', *Plant Cell*, vol. 6, no. 5, pp. 669-683.
- Weinl, S & Kudla, J 2009, 'The CBL-CIPK  $\text{Ca}^{2+}$ -decoding signaling network: Function and perspectives', *New Phytologist*, vol. 184, no. 3, pp. 517-528.
- Wu, C, Gao, X, Kong, X, Zhao, Y & Zhang, H 2009, 'Molecular cloning and functional analysis of a  $\text{Na}^+/\text{H}^+$  antiporter gene *ThNHX1* from a halophytic plant *Thellungiella halophila*', *Plant Molecular Biology Reporter*, vol. 27, no. 1, pp. 1-12.
- Wu, CA, Yang, GD, Meng, QW & Zheng, CC 2004, 'The cotton *GhNHX1* gene encoding a novel putative tonoplast  $\text{Na}^+/\text{H}^+$  antiporter plays an important role in salt stress', *Plant and Cell Physiology*, vol. 45, no. 5, pp. 600-607.
- Wu, H, Shabala, L, Liu, X, Azzarello, E, Zhou, M, Pandolfi, C, Chen, ZH, Bose, J, Mancuso, S & Shabala, S 2015, 'Linking salinity stress tolerance with tissue-specific  $\text{Na}^+$  sequestration in wheat roots', *Frontiers in Plant Science*, vol. 6, no. FEB, pp. 1-13.

- Wu, H, Zhu, M, Shabala, L, Zhou, M & Shabala, S 2014, 'K<sup>+</sup> retention in leaf mesophyll, an overlooked component of salinity tolerance mechanism: A case study for barley', *Journal of Integrative Plant Biology*, pp. 1-15.
- Wu, HH, Shabala, L, Barry, K, Zhou, MX & Shabala, S 2013, 'Ability of leaf mesophyll to retain potassium correlates with salinity tolerance in wheat and barley', *Physiologia Plantarum*, vol. 149, no. 4, pp. 515-527.
- Wu, SJ, Ding, L & Zhu, JK 1996, 'SOS1, a genetic locus essential for salt tolerance and potassium acquisition', *Plant Cell*, vol. 8, no. 4, pp. 617-627.
- Wu, YY, Chen, QJ, Chen, M, Chen, J & Wang, XC 2005, 'Salt-tolerant transgenic perennial ryegrass (*Lolium perenne* L.) obtained by *Agrobacterium tumefaciens*-mediated transformation of the vacuolar Na<sup>+</sup>/H<sup>+</sup> antiporter gene', *Plant Science*, vol. 169, no. 1, pp. 65-73.
- Wyn Jones, RG, Brady, CJ & Speirs, J 1979, *Recent advances in cereal biochemistry*.
- Xu, CX, Zheng, L, Gao, CQ, Wang, C, Liu, GF, Jiang, J & Wang, YC 2011, 'Overexpression of a Vacuolar H<sup>+</sup>-ATPase c subunit gene mediates physiological changes leading to enhanced salt tolerance in transgenic tobacco', *Plant Molecular Biology Reporter*, vol. 29, no. 2, pp. 424-430.
- Xue, ZY, Zhi, DY, Xue, GP, Zhang, H, Zhao, YX & Xia, GM 2004, 'Enhanced salt tolerance of transgenic wheat (*Triticum aestivum* L.) expressing a vacuolar Na<sup>+</sup>/H<sup>+</sup> antiporter gene with improved grain yields in saline soils in the field and a reduced level of leaf Na<sup>+</sup>', *Plant Science*, vol. 167, no. 4, pp. 849-859.
- Yamaguchi, T, Aharon, GS, Sottosanto, JB & Blumwald, E 2005, 'Vacuolar Na<sup>+</sup>/H<sup>+</sup> antiporter cation selectivity is regulated by calmodulin from within the vacuole in a Ca<sup>2+</sup>- and pH-dependent manner', *Proceedings of the National Academy of Sciences of the United States of America*, vol. 102, no. 44, pp. 16107-16112.
- Yeo, AR, Flowers, SA, Rao, G, Welfare, K, Senanayake, N & Flowers, TJ 1999, 'Silicon reduces sodium uptake in rice (*Oryza sativa* L.) in saline conditions and this is accounted for by a reduction in the transpirational bypass flow', *Plant Cell and Environment*, vol. 22, no. 5, pp. 559-565.
- Yeo, AR, Lee, KS, Izard, P, Boursier, PJ & Flowers, TJ 1991, 'Short-term and long-term effects of salinity on leaf growth in rice (*Oryza sativa* L.)', *Journal of Experimental Botany*, vol. 42, no. 240, pp. 881-889.
- Yokoi, S, Quintero, FJ, Cubero, B, Ruiz, MT, Bressan, RA, Hasegawa, PM & Pardo, JM 2002, 'Differential expression and function of *Arabidopsis thaliana* NHX Na<sup>+</sup>/H<sup>+</sup> antiporters in the salt stress response', *Plant Journal*, vol. 30, no. 5, pp. 529-539.
- Yue, YS, Zhang, MC, Zhang, JC, Duan, LS & Li, ZH 2012, 'SOS1 gene overexpression increased salt tolerance in transgenic tobacco by maintaining a higher K<sup>+</sup>/Na<sup>+</sup> ratio', *Journal of Plant Physiology*, vol. 169, no. 3, pp. 255-261.
- Zhang, HX & Blumwald, E 2001a, 'Transgenic salt-tolerant tomato plants accumulate salt in foliage but not in fruit', *Nature Biotechnology*, vol. 19, no. 8, pp. 765-768.

- Zhang, HX, Hodson, JN, Williams, JP & Blumwald, E 2001b, 'Engineering salt-tolerant Brassica plants: Characterization of yield and seed oil quality in transgenic plants with increased vacuolar sodium accumulation', *Proceedings of the National Academy of Sciences of the United States of America*, vol. 98, no. 22, pp. 12832-12836.
- Zhang, HY, Niu, XL, Liu, J, Xiao, FM, Cao, SQ & Liu, YS 2013, 'RNAi-directed downregulation of vacuolar H<sup>+</sup>-ATPase subunit A results in enhanced stomatal aperture and density in rice', *Plos One*, vol. 8, no. 7.
- Zhao, FY, Zhang, XJ, Li, PH, Zhao, YX & Zhang, H 2006, 'Co-expression of the *Suaeda salsa* *SsNHXI* and Arabidopsis *AVPI* confer greater salt tolerance to transgenic rice than the single *SsNHXI*', *Molecular Breeding*, vol. 17, no. 4, pp. 341-353.
- Zhao, Q, Zhao, YJ, Zhao, BC, Ge, RC, Li, M, Shen, YZ & Huang, ZJ 2009, 'Cloning and functional analysis of wheat V-H<sup>+</sup>-ATPase subunit genes', *Plant Molecular Biology*, vol. 69, no. 1-2, pp. 33-46.
- Zhen, RG, Kim, EJ & Rea, PA 1997, 'The molecular and biochemical basis of pyrophosphate-energized proton translocation at the vacuolar membrane', in RA Leigh & D Sanders (eds), *Advances in Botanical Research Incorporating Advances in Plant Pathology, Vol 25: The Plant Vacuole*, vol. 25, pp. 297-337.
- Zhou, GF, Delhaize, E, Zhou, MX & Ryan, PR 2013, 'The barley *MATE* gene, *HvAACT1*, increases citrate efflux and Al<sup>+3</sup> tolerance when expressed in wheat and barley', *Annals of Botany*, vol. 112, no. 3, pp. 603-612.
- Zhou, GF, Delhaize, E, Zhou, MX & Ryan, PR 2014, 'Enhancing the aluminium tolerance of barley by expressing the citrate transporter genes *SbMATE* and *Frd3*', *Journal of Experimental Botany*, vol. 65, pp. 2381-2390.
- Zhou, SF, Chen, XY, Zhang, XG & Li, YX 2008, 'Improved salt tolerance in tobacco plants by co-transformation of a betaine synthesis gene *BADH* and a vacuolar Na<sup>+</sup>/H<sup>+</sup> antiporter gene *SeNHXI*', *Biotechnology Letters*, vol. 30, no. 2, pp. 369-376.
- Zhou, SF, Zhang, ZM, Tang, QL, Lan, H, Li, YX & Luo, P 2011, 'Enhanced V-ATPase activity contributes to the improved salt tolerance of transgenic tobacco plants overexpressing vacuolar Na<sup>+</sup>/H<sup>+</sup> antiporter *AtNHXI*', *Biotechnology Letters*, vol. 33, no. 2, pp. 375-380.
- Zhu, JK 2000, 'Genetic analysis of plant salt tolerance using arabidopsis', *Plant Physiology*, vol. 124, no. 3, pp. 941-948.
- Zhu, JK 2003, 'Regulation of ion homeostasis under salt stress', *Current Opinion in Plant Biology*, vol. 6, no. 5, pp. 441-445.

## Appendix

Cloning of three genes was performed namely Arabidopsis V-ATPase subunit C and subunit D. Mostly, the general steps for developing transgenic lines that the salt group in ACPFG follows are: (1) identifying the Gene of Interest (GOI); (2) searching the gene from the sequence database; (3) designing primers; (4) optimizing PCR condition to amplify the GOI; (5) Inserting the GOI into entry clone-the one used is TOPO gateway entry clone; (6) checking the insert is in correct orientation by restriction digestion; (7) checking the sequence integrity-to avoid SNPs; (8) transferring the GOI to destination Vector (pMDC32) and finally, (9) submitting the GOI in the destination vector for transformation. In addition to this, characterization of Arabidopsis *NHX1* and V-ATPase subunit C genes in barley were made.

### **Generation of overexpression barley lines of V-ATPase subunit C&D**

The Gene sequences were obtained from databases (e.g NCBI) and primers from the cDNA of these sequences were designed. These candidate genes were amplified using PCR and to get a clear band, the PCR condition has to be optimized. Fortunately, it was possible to amplify the V-ATPase subunit C&D using non proof reading DNA polymerase, as using a proof reading, high fidelity polymerase is expensive during optimization step.

After obtaining a clear band of the two V-ATPase genes using normal DNA polymerase, these same genes were amplified using a high fidelity and proof reading capable DNA polymerase called Elongase enzyme mix. The expected band size for subunit D is **786bp** and for subunit C **1128bp**. The bands are presented in the figure below (Fig 1). The DNA ladder used was 100bp. Half of the volume of these PCR products was loaded on to gel and the remaining half purified by PCR purification kit and used further in the cloning process.

The PCR condition to get the bands shown below was as follows:

<u>Components</u>	<u>Volume/reaction</u>
10mM dNTP mix	0.5 $\mu$ L
5X Buffer A	2.5 $\mu$ L
5X Buffer B	2.5 $\mu$ L
Elongase Enzyme mix	0.5 $\mu$ L
<u>MilliQ Water</u>	<u>16.5 <math>\mu</math>L</u>
	22.5 $\mu$ L

#### To the Tubes

Template (root At cDNA)	0.5 $\mu$ L
Forward Primer (10 $\mu$ M)	1 $\mu$ L
<u>Reverse Primer (10<math>\mu</math>M)</u>	<u>1<math>\mu</math>L</u>
	25 $\mu$ L

The thermocycler was programmed as initial denaturation at 94<sup>0</sup>C for 30 seconds, denaturation at 94<sup>0</sup>C for 30 seconds, annealing at 55<sup>0</sup>C for 30 seconds and extension at 68<sup>0</sup>C for 60 seconds. The cycle of denaturation to extension was repeated for 35 cycles.



**Figure 1.** The amplified PCR product of At V-ATPase subunit C and D

As mentioned above, the remaining purified PCR products of the two genes were cloned into pCR8/GW/TOPO TA Gateway @ entry clone (entry vector in short). For doing so, the

following procedure was used. 4  $\mu\text{L}$  of purified PCR product, 1  $\mu\text{L}$  salt solution, 0.5  $\mu\text{L}$  pCR8/GW/TOPO TA Gateway @ entry clone and 0.5  $\mu\text{L}$  of milliQ water were gently mixed and incubated at room temperature for 3 hours covered in aluminium foil. This procedure will allow the GOI to be inserted into entry clone. The next step was to transform the entry clone into competent bacteria cells.

For heat shock transformation, 20  $\mu\text{L}$  TOPO competent cells was thawed on ice for 10-15 minutes and 2  $\mu\text{L}$  of the entry vector containing GOI (pCR8) was added into the competent cells. These mixtures were heat shock for 90 seconds at 42  $^{\circ}\text{C}$  and immediately put back on ice for 5 minutes and 800 $\mu\text{L}$  of SOB solution (to recover the cells) were added in the laminar flow. The above mixture was incubated at 37 $^{\circ}\text{C}$  for 1 hour shaking. After 1 hour, the transformed cells were plated onto a petri-plate containing LB agar with spectinomycines for growing only the transformed cells. The petri –plates were incubated at 37 $^{\circ}\text{C}$  overnight.

From the Petri plates, single colonies were picked by new tips and inoculated LB media and cultured overnight at 37 $^{\circ}\text{C}$  in the incubator-shaker. Finally, Plasmid DNA was extracted from the cultured bacteria cells using Plasmid mini prep kit.

To check the insertion of the gene in the vector, which might be optional to do, restriction digestion using EcoRI was made. Among the 12 colonies of V-ATPase subunit D (designated here as D) and subunit C (designated as C), 11 showed insertion for D and 6 for C, which are D1,D2,D3,D4,D5,D6,D7,D9,D10,D11,D12 & C1,C2,C3,C4,C6,C7 (Fig 2).






Figure 2. EcoRI digestion to see the presence of the GOI in our clones







**Figure 3.** The construct map of VATPase subunit C & D in the entry clone.

### Checking the correct orientation of the GOI in the entry clone

For checking the correct orientation of the GOI in the entry clone, the clone was double digested with two restriction enzymes. HpaI and XhoI for VATPase subunit D gene which gives an expected band size of 1925bp, 1241bp, and 437bp. And, ClaI and PvuII was used for V-ATPase subunit C gene that is expected to give band size of 2188bp, 1186bp, and 571bp.


**Restriction digestion conditions for th VATPase subunit D gene insert**

<u>Components</u>	<u>Volume/reaction</u>
10X NEBuffer	1.0μL
Hpa I	0.2 μL
Xho I	0.2 μL
BSA (100X)	0.1μL
<u>MilliQ Water</u>	<u>6.5 μL</u>
	8.0μL
<u>To the Tubes</u>	
<u>Plasmid DNA</u>	<u>2.0μL</u>
	10μL

**Restriction digestion conditions for th VATPase subunit C gene insert**

<u>Components</u>	<u>Volume/reaction</u>
10X NEBuffer	1.0μL
Cla I	0.2 μL
Pvu II	0.2 μL
BSA (100X)	0.1μL
<u>MilliQ Water</u>	<u>6.5 μL</u>
	8.0μL
<u>To the Tubes</u>	
<u>Plasmid DNA</u>	<u>2.0μL</u>
	10μL


Firstly, out of the 12 clones for each of the genes, 4 clones from each gene were double digested to see whether the gene is correctly oriented in the clone and it was found out that 3 of the clones for the VATPase D gene (D1, D2 and D3) were inserted in the correct orientation. However, only one clone for the VATPase C gene were found to have correct orientation of the insert. The DNA ladder used was 1kb and the clones with correct orientation are marked in bold at the gel picture below (Fig.4).



**Figure 4.** Four clones from each of the D and C genes double digested with restriction enzyme.

The next step, after checking the correct orientation of the gene in the entry clone, is to sequence the gene for examining the integrity of the sequence, i.e., the presence of Single Nucleotide Polymorphisms (SNPs) in the sequence that tend to happen due to the errors introduced to the sequence in all the cloning steps involved. Usually, it is good to sequence more than one clone to widen the chance of getting clones with no SNPs. Nevertheless, in our result explained above, only one clone for the C gene showed correct orientation of the GOI. Hence, additional clones for the C gene have to be tested for correct orientation of the GOI to have at least two clones with correct orientation of the GOI which can be subjected to sequencing. Accordingly, all the available clones of the C gene which had the GOI in them were double digested with their respective restriction enzymes. Among the tested clones, only

one was found to have the GOI in the correct orientation which was the C7 clone. This makes the clones from the C gene two that are ready for sequencing. The C7 clone that showed the correct orientation of the C gene is marked in bold in the gel below (Fig. 5).



**Figure 5.** Three clones from the C gene double digested with restriction enzyme.

### Sequencing to check the presence of SNPs

As described above, it is good to sequence more than one clone and hence, two clones from each gene were sequenced using sense primer GW1 and antisense primer GW2. The sequencing reactions were made three times for each primer per clone which makes it six reactions per clone and a total of 12 per gene and 24 for both genes. The sequencing reaction used the following reagents and PCR conditions.

<u>Components</u>	<u>Volume/reaction</u>
Big Dye Terminator V.3.1	1.0 $\mu$ L
5X Big Dye Terminator dilution buffer	3.5 $\mu$ L
<u>MilliQ Water</u>	<u>4.68 <math>\mu</math>L</u>
	9.18 $\mu$ L
<u>To the Tubes</u>	
Plasmid DNA	0.5 $\mu$ L
<u>Primer (GW1 or GW2)</u>	<u>0.32 <math>\mu</math>L</u>
	10 $\mu$ L

The PCR condition was initial denaturation at 96<sup>0</sup>C for 2 minutes, denaturation at 96<sup>0</sup>C for 10 seconds, annealing at 50<sup>0</sup>C for 5 seconds, extension at 60<sup>0</sup>C for 4 minutes and the cycle is repeated 30 times. Then, to clean up the sequencing reaction, it was allowed to get to room temperature and transferred to 1.5 ml eppendorf tubes. 75 µL of 0.2 mM MgSO<sub>4</sub> in 70 % ethanol was added, mixed by vortexing and allowed to sit at room temperature for 15 minutes. The solution was centrifuged for 15 minutes at 15,000g and the supernatant was removed with a pipette. 75 µL of 70% ethanol was added and centrifuged for 15 minutes at 15,000g. The supernatant removed and the pellet air dried by protecting it from light as long as it dries and submitted for sequencing.

The sequencing result showed that the D gene had SNPs and therefore, the whole process of cloning has to be repeated for this gene. Fortunately, for the C gene, the sequences made using GW1 primers were good except some poor sequencing reads for the GW2 sequences. Hence, the C sequence was repeated only using GW2 primer along with other clones of D gene. Finally, the sequence of the C gene showed no SNPs for the sequence reads in both the sense (GW1) and antisense (GW2) primers that makes it ready to go to the next steps of cloning ,i.e., cloning into the destination vector using LR reaction. The LR reaction components and the volume used and the incubation condition were as follows:

<u>Components</u>	<u>Volume in µL</u>
VATPase C in PCR8	4
pMDC32 (destination vector)	1
5X LR enzyme mix	2
<u>TE Buffer (pH 8)</u>	<u>3</u>
	10


The above solution was incubated at 20<sup>0</sup>C overnight which resulted in the VATPase C gene cloned into the destination vector. Similarly, the C gene cloned in destination vector was

transformed into DH5 $\alpha$  competent bacterial cells following the same protocol described above for the transformation of entry clone. Also, after transformation culturing of the transformed bacterial colonies was done similar to the entry clone. Following culturing of the transformed bacterial cells, the plasmid was extracted using plasmid mini prep kit (uses resuspension, lysis and neutralization solutions).

To check the presence of the VATPase C gene in the pMDC32 restriction enzyme EcoRI was selected and the banding pattern was determined with Vector NTI software. The banding pattern expected from this restriction enzyme was 9761bp, 1146bp and 345bp. The volume of the components and the incubation condition is presented below.

<u>Components</u>	<u>Volume/reaction</u>
EcoR I (promega)	0.2 $\mu$ L
Promega EcoR I buffer (H)	1.5 $\mu$ L
<u>MilliQ water</u>	<u>11.3 <math>\mu</math>L</u>
	13 $\mu$ L
<u>DNA</u>	<u>2 <math>\mu</math>L</u>
	15 $\mu$ L


The above solution was incubated overnight at 37<sup>0</sup>C. Then, the digested DNA was run on agarose gel and the correct banding pattern was observed in the first clone which is shown in the first lane of figure below (Fig. 6)



**Figure 6.** The restrictions digest of the pMDC32 to check the presence of VATPase C gene.

To further confirm that this clone indeed contain the VATPase gene, another restriction enzyme (Pvu I) was used. Similarly, the banding pattern to be generated by using this restriction enzyme was determined using Vector NTI software. It was 6671bp, 2719bp, 1675bp and 687bp. Consistent to the above result, the same clone showed the expected banding pattern confirming that it contains the VATPase C gene in the pMDC32 (Fig. 7)

All the below clones are derived from C7 clone in pMDC32 (destination vector)




**Figure 7.** The Pvu I restriction enzyme digestion to further confirm the presence of VATPase C gene in pMDC32.

Finally, this clone that contains our gene was sent to the transformation lab for barley transformation.

### **Repeating the cloning of VATPase subunit D gene**


Because the sequencing result for VATPase D gene had SNPs, the process of cloning has been done again following the steps described above. The gel picture showed the successful amplification of the gene using the same protocol described above. All the three lanes in the figure shows the same gene amplified in three tubes with similar condition (Fig. 8)



**Figure 8.** VATPase subunit D gene amplified using elongase enzyme

The gene was cloned into pCR8 entry clone and digested for correct orientation similar to the previous condition and also with similar enzymes. The double digestion was made using HpaI and XhoI and the expected band sizes were 1925bp, 1241bp, and 437bp. The D11 & D12 clones showed the correct orientation (Fig. 9). One of the clones which showed the correct orientation will be cloned into pMDC32.





**Figure 9.** Double digestion to check correct orientation of V-ATPase D inserts.

### **Peptide sample preparation and analysis by nano liquid chromatography/ Tandem mass spectrometry**

Peptide samples were generated using co-precipitation of proteins with trypsin as previously described (Wilson et al., 2010). Briefly, 100  $\mu$ l aliquots of crude protein extract were clarified by centrifugation at 13,000 rpm and the supernatant precipitated with 9 volumes of ice-cold ethanol (2 hrs at -20). The protein pellets were resuspended in 90  $\mu$ l denaturing buffer (7M urea, 2M thiourea in 30mM Tris pH 8.0) and subjected to sequential reduction, alkylation and trypsin digestion.

Peptide samples were analysed by nanoLC-MS/MS using an LTQ-Orbitrap XL (ThermoFisher Scientific, Waltham, MA, USA). Aliquots of tryptic peptides were loaded at 0.05 ml/min onto a C18 capillary trapping column (Peptide CapTrap, Michrom BioResources, Auburn, CA, USA) controlled by an Alliance 2690 Separations Module (Waters). Peptides

were then separated using a Surveyor MS Pump plus (ThermoFisher Scientific) on an analytical nanoHPLC column packed with 5 micron C18 ProteoPep II media (PicoFrit Column, 15  $\mu$ m i.d. pulled tip, 10 cm, New Objective) as previously described (Al-Naseri *et al.* 2013).

The LTQ-Orbitrap XL was controlled using Xcalibur 2.0 software (ThermoFisher Scientific) and operated in data-dependent acquisition mode where survey scans were acquired in the Orbitrap using a resolving power of 60,000 (at 400 m/z). MS/MS spectra were concurrently acquired in the LTQ mass analyser on the eight most intense ions from the FT survey scan. Charge state filtering, where unassigned and singly-charged precursor ions were not selected for fragmentation, and dynamic exclusion (repeat count 1, repeat duration 30 s, exclusion list size 500) were used. Fragmentation conditions in the LTQ were: 35% normalised collision energy, activation q of 0.25, 30 ms activation time and minimum ion selection intensity of 500 counts.

### **Database searching and criteria for protein identification**

Centroid mode spectra acquired were converted from .RAW files into .mzXML peak list files using the msConvert command (Proteowizard). The extracted MS/MS data were searched against the Barley protein database of 36,116 entries downloaded from the National Centre for Biotechnology Information on 21/05/2013, to which sequence for the Arabidopsis sodium hydrogen exchange protein -1 had been appended (>sp|Q68KI4|NHX1\_ARATH Sodium/hydrogen exchanger 1 OS=Arabidopsis thaliana GN=NHX1). Semi-tryptic searches using parent ion and fragment ion mass tolerances of 10 ppm and 0.5 Da, respectively, were performed using X!Tandem running in the Computational Proteomics Analysis System (CPAS), an open-source bioinformatics resource for analysing large proteomics datasets (Rauch *et al.* 2006).

S-carboxamidomethylation of cysteine residues was specified as a fixed modification and oxidation of methionine was specified as a variable modification. The Peptide Prophet and Protein Prophet algorithms were applied to the X!Tandem search results to assign probabilities to peptide and protein matches, respectively (Keller *et al.* 2002; Nesvizhskii *et al.* 2003). Peptide-spectrum matches were accepted if the peptide was assigned a probability > 0.95 by the Peptide Prophet algorithm. Protein identifications were accepted if the protein contained two or more unique peptide sequences and the protein was assigned a probability > 0.99 by the Protein Prophet algorithm.


Al-Naseri, A, Bowman, JP, Wilson, R, Nilsson, RE, & Britz, ML 2013,' Impact of Lactose Starvation on the Physiology of *Lactobacillus casei* GCRL163 in the Presence or Absence of Tween 80', *Journal of Proteome Research*, Vol. 12, no. 11, pp 5313-5322.

Rauch, A, Bellew, M, Eng, J, Fitzgibbon, M, Holzman, T, Hussey, P, Igra, M, Maclean, B, Lin, WC, Detter, A, Fang, R, Faca, V, Gafken, P, Zhang, H, Whitaker, J, States, D, Hanash, S, Paulovich, A, & Macintosh, MW 2006, 'Computational Proteomics Analysis System (CPAS): An Extensible, Open-Source Analytic System for Evaluating and Publishing Proteomic Data and High Throughput Biological Experiments', *Journal of Proteome Research*, vol. 5, no.1, pp. 112-121.


Keller, A, Nesvizhskii, AI, Kolker, E, & Aebersold, R 2002, 'Empirical statistical model to estimate the accuracy of peptide identifications made by MS/MS and database search', *Analytical Chemistry*, vol.74, pp. 5383-92.

Nesvizhskii, AI, Keller, A, Kolker, E, & Aebersold, R 2003, 'A Statistical Model for Identifying Proteins by Tandem Mass Spectrometry', *Analytical Chemistry*, vol. 75, pp. 4646-58.

Wilson, R, Diseberg, AF, Goordon, L, Zivkovic,S, Tatarczuch, L, Mackie, EJ, Gorman, JJ & Bateman, JF 2010,' Comprehensive profiling of cartilage extracellular matrix formation & maturation using sequential extraction & label-free quantitative proteomics', *Molecular & Cellular Proteomics*, vol. 9, issue 6, pp. 1296-1313.



1. Sample ID #10734 ion map view (intensity scaled by colour – green is most intense signal)
2. Example high-resolution FTMS survey scan (resolving power 60,000)



3. Sample ID #10734 base peak chromatogram

4. Example MS/MS spectrum for precursor ion m/z 909.99 – see details on following slides

gi|112684|sp|P29305.1|1433A\_HORVU (1433A\_HORVU)

Sequence Mass:29,352

AA Coverage:56% (146 / 262)

Mass Coverage:55% (16,139 / 29,352)

MSTAEATREE NVYMAKLAEQ AERYEEMVEF MEKVAKTADV GELTVEERNL LSVAYKNVIG ARRASWRIIS SIEQKEESRG NEAYVASIKE YRTRITETLS KICDGILKLL DSHLVPSATA AESKVFYLMK KGDYHRYLAE FKAGAERKEA AENTLVAYKS AQDIALADLP TTHPIRLGLA LNFVVFYIEI LNSPDRACNL AKQAFDEAIA ELDLGEESY KDSTLIMQLL RDNLTILWTS NAEFGGDEIK EAASKPEGEG HS

Annotations for 1433A\_HORVU

Sequence name: 1433A\_HORVU

Description: 14-3-3-like protein A - Hordeum vulgare (Barley)

Gene Name(s):

Organisms(s): Hordeum vulgare

Genbank IDs

GIs

Swiss-Prot Accessions

Swiss-Prot Names

Ensembl

IPI numbers

GO Categories

112684

P29305.1

1433A\_HORVU


P29305

Peptides

Peptide Filter: (PeptideProphet >= 0.95)

Peptide Sort: Protein ASC, Peptide ASC

Scan	Z	Hyper	Next	B	Y	Expect	Ion%	CalcMH+	dMass	PepProphet	Peptide	SeqHits
4379	2+	46.8	34.2	10.5	11.5	0.5300	83%	1189.6609	+0.0050	0.9937	K.DSTLIMQLLR.D	27
4454	2+	46.9	37.7	10.7	11.5	0.2700	83%	1189.6609	+0.0050	0.9934	K.DSTLIMQLLR.D	27
1490	2+	47.6	45.2	7.4	10.2	0.0360	80%	1208.6157	+0.0040	0.9946	K.EAAENTLVAYK.S	3
1900	2+	72.3	22.9	11.9	13.0	0.0000	70%	1638.8697	+0.0040	1.0000	K.LLDShLVPSATAAESK.V	3
10325	3+	46.7	34.2	11.6	11.0	0.0056	17%	3268.5930	+0.0170	0.9983	K.Q^AFDEAIAELDSLGEESYKDSTLIMQLLR.D	3
5311	2+	67.7	34.5	9.6	12.2	0.0000	56%	2114.9764	+0.0090	1.0000	K.QAFDEAIAELDSLGEESYK.D	3
5350	3+	49.1	45.5	10.0	10.5	0.3600	29%	2114.9760	+0.0150	0.9753	K.QAFDEAIAELDSLGEESYK.D	3
5401	2+	49.8	27.4	8.8	11.8	0.0007	44%	2114.9764	-0.0010	0.9997	K.QAFDEAIAELDSLGEESYK.D	3
2510	2+	79.2	35.1	10.8	13.5	0.0000	72%	1818.9708	+0.0050	1.0000	K.SAQDIALADLPTTHPIR.L	3
2514	3+	61.1	37.1	10.0	12.4	0.0013	38%	1818.9708	+0.0040	1.0000	K.SAQDIALADLPTTHPIR.L	3
1667	2+	45.9	40.0	8.7	10.9	0.0500	68%	1318.6485	+0.0070	0.9967	K.TADVGLTVEER.N	3
2836	3+	107.9	46.9	7.6	11.1	0.0000	32%	3199.4298	+0.0050	1.0000	R.DNLTILWTSNNAEEGGDEIKEAASKPEGEGH.S	3
1019	2+	73.9	35.8	8.8	12.3	0.0000	100%	1418.7485	+0.0050	1.0000	R.IISSIEQKEESR.G	12
860	2+	41.6	30.2	11.4	9.8	0.2400	88%	1076.5946	+0.0030	0.9654	R.TRIETELSK.I	3
2518	2+	57.3	28.6	10.1	12.2	0.0002	100%	1334.5643	+0.0060	1.0000	R.YEEMVEFMEK.V	12



$b^{2+}$	$b^{+}$	#	AA	#	$y^{+}$	$y^{2+}$
44.5470	88.0861	1	S	17		
80.0864	159.1649	2	A	16	1732.9774	866.9927
144.1518	287.2957	3	Q	15	1661.8986	831.4533
201.6961	402.3843	4	D	14	1533.7679	767.3879
258.2758	515.5437	5	I	13	1418.6793	709.8436
293.8152	586.6225	6	A	12	1305.5198	653.2639
350.3949	699.7819	7	L	11	1234.4410	617.7245
385.9343	770.8607	8	A	10	1121.2816	561.1448
443.4786	885.9493	9	D	9	1050.2028	525.6054
500.0584	999.1088	10	L	8	935.1142	468.0611
548.6167	1096.2255	11	P	7	821.9547	411.4813
599.1692	1197.3305	12	T	6	724.8381	362.9230
649.7218	1298.4356	13	T	5	623.7330	312.3705
718.2923	1435.5767	14	H	4	522.6279	261.8179
766.8507	1532.6934	15	P	3	385.4868	193.2474
823.4304	1645.8528	16	I	2	288.3701	144.6890
		17	R	1	175.2107	88.1093

MS/MS spectrum (daughter ions of peptide m/z 909)





